

Subcritical phase of d -dimensional Poisson-Boolean percolation and its vacant set

Hugo Duminil-Copin*, Aran Raoufi†, Vincent Tassion‡

May 13, 2019

Abstract

We prove that the Poisson-Boolean percolation on \mathbb{R}^d undergoes a sharp phase transition in any dimension under the assumption that the radius distribution has a $5d - 3$ finite moment (in particular we do not assume that the distribution is bounded). To the best of our knowledge, this is the first proof of sharpness for a model in dimension $d \geq 3$ that does not exhibit exponential decay of connectivity probabilities in the subcritical regime. More precisely, we prove that in the whole subcritical regime, the expected size of the cluster of the origin is finite, and furthermore we obtain bounds for the origin to be connected to distance n : when the radius distribution has a finite exponential moment, the probability decays exponentially fast in n , and when the radius distribution has heavy tails, the probability is equivalent to the probability that the origin is covered by a ball going to distance n (this result is new even in two dimensions). In the supercritical regime, it is proved that the probability of the origin being connected to infinity satisfies a mean-field lower bound. The same proof carries on to conclude that the vacant set of Poisson-Boolean percolation on \mathbb{R}^d undergoes a sharp phase transition.

1 Introduction

Definition of the model Bernoulli percolation was introduced in [BH57] by Broadbent and Hammersley to model the diffusion of a liquid in a porous medium. Originally defined on a lattice, the model was later generalized to a number of other contexts. Of particular interest is the development of continuum percolation (see [MR08] for a book on the subject), whose most classical example is provided by the Poisson-Boolean model (introduced by Gilbert [Gil61]). It is defined as follows.

Fix a positive integer $d \geq 2$ and let \mathbb{R}^d be the d -dimensional Euclidean space endowed with the ℓ^2 norm $\|\cdot\|$. For $r > 0$ and $x \in \mathbb{R}^d$, set $B_r^x := \{y \in \mathbb{R}^d : \|y - x\| \leq r\}$ and

*Institut des Hautes Études Scientifiques and Université de Genève, duminil@ihes.fr

†ETH Zürich, aran.raoufi@math.ethz.ch

‡ETH Zürich, vincent.tassion@math.ethz.ch

$\partial\mathbf{B}_r^x := \{y \in \mathbb{R}^d : \|y - x\| = r\}$ for the ball and sphere of radius r centered at x . When $x = 0$, we simply write \mathbf{B}_r and $\partial\mathbf{B}_r$. For a subset η of $\mathbb{R}^d \times \mathbb{R}_+$, set

$$\mathcal{O}(\eta) := \bigcup_{(z,R) \in \eta} \mathbf{B}_R^z.$$

Let μ be a measure on \mathbb{R}_+ (below, we use the notation $\mu[a, b]$ to refer to $\mu([a, b])$, where $a, b \in \mathbb{R} \cup \{\infty\}$) and λ be a positive number. Let η be a Poisson point process of intensity $\lambda \cdot dz \otimes \mu$, where dz is the Lebesgue measure on \mathbb{R}^d . Write \mathbb{P}_λ for the law of η and \mathbb{E}_λ for the expectation with respect to \mathbb{P}_λ . The random variable $\mathcal{O}(\eta)$, where η has law \mathbb{P}_λ is called the *Poisson-Boolean percolation* of radius law μ and intensity λ . A natural hypothesis in the study of Poisson Boolean percolation is to assume d -th moment on the radius distribution:

$$\int_0^\infty r^d d\mu(r) < \infty. \quad (1)$$

Indeed, as observed by Hall [Hal85], the condition (1) is necessary in order to avoid the entire space to be almost surely covered, regardless of the intensity (as long as positive) of the Poisson point process.

Main result Two points x and y of \mathbb{R}^d are said to be *connected* (by η) if there exists a continuous path in $\mathcal{O}(\eta)$ connecting x to y . This event is denoted by $x \longleftrightarrow y$. For $X, Y \subset \mathbb{R}^d$, the event $\{X \longleftrightarrow Y\}$ denotes the existence of $x \in X$ and $y \in Y$ such that x is connected to y (when $X = \{x\}$, we simply write $x \longleftrightarrow Y$). Define, for every $r > 0$, the two functions of λ

$$\theta_r(\lambda) := \mathbb{P}_\lambda[0 \longleftrightarrow \partial\mathbf{B}_r] \quad \text{and} \quad \theta(\lambda) := \lim_{r \rightarrow \infty} \theta_r(\lambda).$$

We define the critical parameter $\lambda_c = \lambda_c(d)$ of the model by the formula

$$\lambda_c := \inf\{\lambda \geq 0 : \theta(\lambda) > 0\}.$$

Another critical parameter is often introduced to discuss Poisson-Boolean percolation. Define $\tilde{\lambda}_c = \tilde{\lambda}_c(d)$ by the formula

$$\tilde{\lambda}_c := \inf\{\lambda \geq 0 : \inf_{r>0} \mathbb{P}_\lambda[\mathbf{B}_r \longleftrightarrow \partial\mathbf{B}_{2r}] > 0\}.$$

This quantity is of great use as it enables one to initialize renormalization arguments; see e.g. [Gou08, GT18] and references therein. As a consequence, a lot is known for Poisson-Boolean percolation with intensity $\lambda < \tilde{\lambda}_c$. We refer to Theorems 2 and 3 and to [MR08] for more details on the subject.

Under the minimal assumption (1), Gouéré [Gou08] proved that λ_c and $\tilde{\lambda}_c$ are non-trivial. More precisely he proved $0 < \tilde{\lambda}_c \leq \lambda_c < \infty$. The main result of this paper is the following.

Theorem 1 (Sharpness for Poisson-Boolean percolation). *Fix $d \geq 2$ and assume that*

$$\int_{\mathbb{R}_+} r^{5d-3} d\mu(r) < \infty. \quad (2)$$

Then, we have that $\lambda_c = \tilde{\lambda}_c$. Furthermore, there exists $c > 0$ such that $\theta(\lambda) \geq c(\lambda - \lambda_c)$ for any $\lambda \geq \lambda_c$.

The case of bounded radius is already proved in [ZS85, MRS94] (see also [MR08]), and we refer the reader to [Zie16] for a new proof. Theorems stating sharpness of phase transitions for percolation models in general dimension d were first proved in the eighties for Bernoulli percolation [Men86, AB87] and the Ising model [ABF87]. The proofs of sharpness for these models (even alternative proofs like [DT16]) harvested independence for Bernoulli percolation and special structures of the random-current representation of the Ising model. In particular, they were not applicable to other models of statistical mechanics. In recent years, new methods were developed to prove sharpness for a large variety of statistical physics models in two dimensions [BR06, BD12, ATT16]. These methods rely on general sharp threshold theorems for Boolean functions, but also on *planar* properties of crossing events. In particular, the proofs use planarity in an essential way and seem impotent in higher dimensions. Recently, the authors proved the sharpness of phase transition for random-cluster model [DRT17b] and Voronoi percolation [DRT17a] in arbitrary dimensions. The shared theme of the proofs is the use of randomized algorithms to prove differential inequalities for connection probabilities.

Here, we do not prove that connectivity probabilities decay exponentially fast in the distance below criticality since this is simply not true in general for Poisson-Boolean percolation. Indeed, if the tail of the radius distribution is slower than exponential, then the event that a large ball covers two given points has a probability larger than exponential in the distance between the two points. Instead, we show by contradiction that $\lambda_c = \tilde{\lambda}_c$, without ever referring to exponential decay, by controlling the derivative of $\theta_r(\lambda)$ in terms of “pivotality events” for λ in the “fictitious” regime $(\tilde{\lambda}_c, \lambda_c)$ and by deriving a differential inequality on $\theta_r(\lambda)$ (using randomized algorithms) which is valid *only* in this regime. The regime $(\tilde{\lambda}_c, \lambda_c)$ is referred to as the *fictitious regime* since one eventually concludes from our proof that the interval is empty.

We wish to highlight that while the proof below also harvests randomized algorithms, we consider that the main novelty of the paper lies in the comparison between the derivative of $\theta_r(\lambda)$ and pivotal events. In many percolation models constructed from disordered systems with long range interaction, the pivotality events that typically appear in Russo’s formula for Bernoulli percolation get replaced by more complicated quantities. For instance, in our model probability of large ball being pivotal also enter into the expression of the derivative, in percolation models built out of Gaussian processes such as Gaussian Free Field or Bargmann-Fock, the probability of pivotality *conditioned on* the field being equal to a certain value are relevant, etc. It is not true that these quantities are comparable to the pivotality probabilities for arbitrary values of λ , but one may show with some work that this is indeed the case in the fictitious regime. We believe that the introduction of this fictitious regime is therefore a powerful tool towards a better

understanding of these models. To illustrate recent applications of similar techniques that were inspired by the present paper, we refer e.g. to [DGRSY18, DGRS19].

Decay of $\theta_r(\lambda)$ when $\lambda < \tilde{\lambda}_c$ For standard percolation, sharpness of the phase transition refers to the exponential decay of connection probabilities in the subcritical regime. In Poisson-Boolean percolation with arbitrary radius law μ , we mentioned above that one cannot expect such behavior to hold in full generality. In order to explain why the theorem above is still called “sharpness” in this article, we provide below some new results concerning the behavior of Poisson-Boolean percolation when $\lambda < \tilde{\lambda}_c$.

First, remark that for every $\lambda > 0$, $\theta_r(\lambda)$ is always bounded from below by

$$\phi_r(\lambda) := \mathbb{P}_\lambda[\exists(z, R) \in \eta \text{ such that } \mathbb{B}_R^z \text{ contains } 0 \text{ and intersects } \partial\mathbb{B}_r], \quad (3)$$

whose decay may be arbitrarily slow. Nevertheless, one may expect the following phenomenology when $\lambda < \tilde{\lambda}_c$:

- If $\mu[r, \infty]$ decays exponentially fast, then so does $\theta_r(\lambda)$ (but not necessarily at the same rate of exponential decay).
- Otherwise, the decay of $\theta_r(\lambda)$ is governed by $\phi_r(\lambda)$, in the sense that it is roughly equivalent to it.

The first item above is formalized by the following theorem.

Theorem 2. *If there exists $c > 0$ such that $\mu[r, \infty] \leq \exp(-cr)$ for every $r \geq 1$, then, for every $\lambda < \tilde{\lambda}_c$, there exists $c_\lambda > 0$ such that for every $r \geq 1$,*

$$\theta_r(\lambda) \leq \exp(-c_\lambda r).$$

Giving sense to the second item above is not easy in full generality, for instance when the law of μ is very irregular (one can imagine distributions μ that do not decay exponentially fast, but for which large range of radii are excluded). In Section 4.3, we give a general condition under which a precise description of $\theta_r(\lambda)$ can be obtained. To avoid introducing technical notation here, we only give two applications of the results proved in Section 4.3. We believe that these applications bring already a good idea of the general phenomenology. The proof mostly relies on new renormalization inequalities. We believe that these renormalization inequalities can be of great use to other percolation models such as long-range Bernoulli percolation models. The theorem claims that the cheapest way for 0 to be connected to distance r is if a single huge ball covers 0 and intersects the boundary of \mathbb{B}_r .

Theorem 3. *Fix $d \geq 2$. Let μ is of one of the two following cases:*

C1 There exists $c > 0$ such that $\mu[r, \infty] = 1/r^{d+c}$ for every $r \geq 1$,

C2 There exist $c > 0$ and $0 < \alpha < 1$ such that $\mu[r, \infty] = \exp(-cr^\alpha)$ for every $r \geq 1$.

Then, for every $\lambda < \tilde{\lambda}_c$,

$$\lim_{r \rightarrow \infty} \frac{\theta_r(\lambda)}{\phi_r(\lambda)} = 1.$$

This theorem is new even in two dimensions. The proofs of Theorem 2 and Theorem 3 are independent of the proof of Theorem 1 and can be read separately.

Vacant set of the Poisson-Boolean model Another model of interest can also be studied using the same techniques. In this model, the connectivity of the points is given by continuous paths in the complement of $\mathcal{O}(\eta)$. Write $x \overset{*}{\longleftrightarrow} y$ for the event that x and y are connected by a continuous path in $\mathbb{R}^d \setminus \mathcal{O}(\eta)$, and $X \overset{*}{\longleftrightarrow} Y$ if there exist $x \in X$ and $y \in Y$ such that x and y are connected. For λ and $r \geq 0$, define

$$\theta_r^*(\lambda) := \mathbb{P}_\lambda[0 \overset{*}{\longleftrightarrow} \partial B_r] \quad \text{and} \quad \theta^*(\lambda) = \lim_{r \rightarrow \infty} \theta_r^*(\lambda).$$

We define the critical parameter λ_c^* (see e.g. [Pen17] or [ATT17] for the fact that it is positive) by the formula

$$\lambda_c^* := \sup\{\lambda \geq 0 : \theta^*(\lambda) > 0\}.$$

We have the following theorem.

Theorem 4 (Sharpness of the vacant set of the Poisson-Boolean model). *Fix $d \geq 2$ and assume that the radius distribution μ is compactly supported. Then, for all $\lambda < \lambda_c^*$, there exists $c_\lambda > 0$ such that for every $r \geq 1$,*

$$\theta_r^*(\lambda) \leq \exp(-c_\lambda r).$$

Furthermore, there exists $c > 0$ such that for every $\lambda \geq \lambda_c^*$, $\theta^*(\lambda) \geq c(\lambda - \lambda_c^*)$.

Since the proof follows the same lines as in Theorem 1, we omit it in the article and leave it as an exercise to the reader.

Strategy of the proof of Theorem 1 Let us now turn to a brief description of the general strategy to prove our main theorem. Theorem 1 is a consequence of the following lemma.

Lemma 5. *Assume the moment condition (2) on the radius distribution. Then, there exists a constant $c_1 > 0$ such that for every $r \geq 0$ and $\lambda \geq \tilde{\lambda}_c$,*

$$\theta_r'(\lambda) \geq c_1 \frac{r}{\Sigma_r(\lambda)} \theta_r(\lambda)(1 - \theta_r(\lambda)), \quad (4)$$

where $\Sigma_r(\lambda) := \int_0^r \theta_s(\lambda) ds$.

The whole point of Section 3 will be to prove Lemma 5. Before that, let us mention how it implies Theorem 1.

Proof of Theorem 1. Fix $\lambda_0 > \tilde{\lambda}_c$. As in [DRT17b, Lemma 3.1], Lemma 5 implies that there exists $\lambda_1 \in [\tilde{\lambda}_c, \lambda_0]$ such that

- for any $\lambda \geq \lambda_1$, $\theta(\lambda) \geq c(\lambda - \lambda_1)$.
- for any $\lambda \in (\tilde{\lambda}_c, \lambda_1)$, there exists $c_\lambda > 0$ such that $\theta_r(\lambda) \leq \exp(-c_\lambda r)$ for every $r \geq 0$.

The two items imply that $\lambda_1 = \lambda_c$. Yet, the second item implies that $\lambda_1 \leq \tilde{\lambda}_c$, since clearly exponential decay would imply that for $\lambda \in (\tilde{\lambda}_c, \lambda_1)$,

$$\lim_{r \rightarrow \infty} \mathbb{P}_\lambda[\mathbf{B}_r \leftrightarrow \partial \mathbf{B}_{2r}] = 0.$$

Since $\lambda_1 = \lambda_c \geq \tilde{\lambda}_c$, we deduce that $\lambda_1 = \lambda_c = \tilde{\lambda}_c$ and the proof is finished. \square

Remark 6. *Note that we did not deduce anything from Lemma 5 about exponential decay since eventually $\lambda_1 = \tilde{\lambda}_c$. It is therefore not contradictory with the case in which $\mu[r, \infty]$ does not decay exponentially fast.*

The proof of Lemma 5 relies on the OSSS inequality, first proved in [OSSS05], connecting randomized algorithms and influences in a product space. Let us briefly describe this inequality. Let I be a finite set of *coordinates*, and let $\Omega = \prod_{i \in I} \Omega_i$ be a product space endowed with product measure $\pi = \otimes_{i \in I} \pi_i$. An *algorithm* T determining $f : \Omega \rightarrow \{0, 1\}$ takes a configuration $\omega = (\omega_i)_{i \in I} \in \Omega$ as an input, and reveals the value of ω in different $i \in I$ one by one. At each step, which coordinate will be revealed next depends on the values of ω revealed so far. The algorithm stops as soon as the value of f is the same no matter the values of ω on the remaining coordinates. Define the functions $\delta_i(\mathsf{T})$ and $\text{Inf}_i(f)$, which are respectively called the *revelment* and the *influence* of the i -th coordinate, by

$$\begin{aligned} \delta_i(\mathsf{T}) &:= \pi[\mathsf{T} \text{ reveals the value of } \omega_i], \\ \text{Inf}_i(f) &:= \pi[f(\omega) \neq f(\tilde{\omega})], \end{aligned}$$

where $\tilde{\omega}$ denotes the random element in Ω^I which is the same as ω in every coordinate except the i -th coordinate which is *resampled independently*.

Theorem 7 ([OSSS05]). *For every function $f : \Omega \rightarrow \{0, 1\}$, and every algorithm T determining f ,*

$$\text{Var}_\pi(f) \leq \sum_{i \in I} \delta_i(\mathsf{T}) \text{Inf}_i(f), \tag{OSSS}$$

where Var_π is the variance with respect to the measure π .

This inequality is used as follows. First, we write Poisson-Boolean percolation as a product space. Second, we exhibit an algorithm for the event $\{0 \leftrightarrow \partial \mathbf{B}_r\}$ for which we control the revelations. Then, we use the assumption $\lambda > \tilde{\lambda}_c$ to connect the influences of the product space to the derivative of θ_r . Altogether, these steps lead to (4).

Organization of the article The next section contains some preliminaries. Section 3 contains the proof of Theorem 1 while the last section contains the proofs of Theorems 2 and 3.

2 Background

We introduce some notation and recall three properties of the Poisson-Boolean percolation that we will need in the proofs of the next sections.

Further notation For $x \in \mathbb{Z}^d$, introduce the squared box $\mathbf{S}^x := x + [-1/2, 1/2]^d$ around x . In order to apply the OSSS inequality, we wish to write our probability space as a product space. To do this, we introduce the following notation. For any integer $n \geq 1$ and $x \in \mathbb{Z}^d$, let

$$\eta_{(x,n)} := \eta \cap (\mathbf{S}^x \times [n-1, n]),$$

which corresponds to all the balls of η centered at a point in \mathbf{S}^x with radius in $[n-1, n)$. All the constants c_i below (in particular in the lemmata) are independent of all the parameters.

Insertion tolerance We will need the following insertion tolerance property. Consider \mathbf{r}_* and \mathbf{r}^* such that

$$\mathbb{P}_\lambda[\mathcal{D}_x] := c_{\text{IT}} = c_{\text{IT}}(\lambda) > 0, \quad (\text{IT})$$

where \mathcal{D}_x is the event that there exists $(z, R) \in \eta$ with $z \in \mathbf{S}^x$ and $\mathbf{B}_{\mathbf{r}_*}^x \subset \mathbf{B}_R^z \subset \mathbf{B}_{\mathbf{r}^*}^x$. Without loss of generality (the radius distribution may be scaled by a constant factor), we further assume that \mathbf{r}_* and \mathbf{r}^* satisfy the following conditions (these will be useful at different stages of the proof):

$$1 + 2\sqrt{d} \leq \mathbf{r}_* \leq \mathbf{r}^* \leq 2\mathbf{r}_* - 2\sqrt{d}. \quad (5)$$

While the quantity c_{IT} varies with λ , the dependency will be continuous and therefore irrelevant for our arguments. We will omit to refer to this subtlety in the proofs to avoid confusion.

FKG inequality An *increasing* event A is an event such that $\eta \in A$ and $\eta \subset \eta'$ implies $\eta' \in A$. The FKG inequality for Poisson point processes states that for every $\lambda > 0$ and every two increasing events A and B ,

$$\mathbb{P}_\lambda[A \cap B] \geq \mathbb{P}_\lambda[A] \mathbb{P}_\lambda[B]. \quad (\text{FKG})$$

Russo's formula For $x \in \mathbb{Z}^d$ and an increasing event A , define the random variable

$$\text{Piv}_{x,A}(\eta) := \mathbf{1}_{\eta \notin A} \int_{\mathbf{S}^x} \int_{\mathbb{R}_+} \mathbf{1}_{\eta \cup (z,r) \in A} dz \mu(dr). \quad (6)$$

Russo's formula yields that

$$\frac{d}{d\lambda} \mathbb{P}_\lambda[A] = \sum_{x \in \mathbb{Z}^d} \mathbb{E}_\lambda[\text{Piv}_{x,A}]. \quad (\text{Russo})$$

3 Proof of Lemma 5

The next subsection contains the proof of Lemma 5 conditioned on two lemmata, which are proved in the next two subsections.

3.1 Proof of Lemma 5

As mentioned above, the proof of the lemma is obtained by applying the OSSS inequality to a truncated version of the probability space generated by the independent variables $(\eta_{(x,n)})_{x \in \mathbb{Z}^d, n \geq 1}$. In this section, we fix $L \geq 2r > 0$. Set $\mathcal{A} := \{0 \longleftrightarrow \partial \mathbb{B}_r\}$ and $f = \mathbf{1}_{\mathcal{A}}$. Define

$$I_L := \{(x, n) \in \mathbb{Z}^d \times \mathbb{N} \text{ such that } \|x\| \leq L \text{ and } 1 \leq n \leq L\}.$$

Also, for $r \geq 0$, let $\eta_{\mathbf{g}}$ denote the union of the $\eta_{(x,n)}$ for $(x, n) \notin I_L$. For i being either (x, n) or \mathbf{g} , set Ω_i to be the space of possible η_i and π_i the law of η_i under \mathbb{P}_λ .

For $0 \leq s \leq r$, apply the OSSS inequality (Theorem 7) to

- the product space $(\prod_{i \in I} \Omega_i, \otimes_{i \in I} \pi_i)$ where $I := I_L \cup \{\mathbf{g}\}$,
- the indicator function $f = \mathbf{1}_{\mathcal{A}}$ considered as a function from $\prod_{i \in I} \Omega_i$ onto $\{0, 1\}$,
- the algorithm $\mathbb{T}_{s,L}$ defined below.

Definition 8 (Algorithm $\mathbb{T}_{s,L}$). *Fix a deterministic ordering of I . Set $i_0 = \mathbf{g}$, and reveal $\eta_{\mathbf{g}}$. At each step t , assume that $\{i_0, \dots, i_{t-1}\} \subset I$ has been revealed. Then,*

- *If there exists $(x, n) \in I \setminus \{i_0, \dots, i_{t-1}\}$ such that the Euclidean distance between \mathbb{S}^x and the connected components of $\partial \mathbb{B}_s$ in $\mathcal{O}(\eta_{i_1} \cup \dots \cup \eta_{i_{t-1}})$ is smaller than n , then set i_{t+1} to be the smallest such (x, n) for some fixed ordering.*
- *If such an (x, n) does not exist, halt the algorithm.*

Remark 9. *Roughly speaking, the algorithm checks $\mathcal{O}(\eta_{\mathbf{g}})$ and then discovers the connected components of $\partial \mathbb{B}_s$.*

Theorem 7 implies that

$$\theta_r(1 - \theta_r) \leq 2 \sum_{i \in I} \delta_i(\mathbb{T}_{s,L}) \text{Inf}_i(f). \quad (7)$$

By construction, the random variable $\eta_{\mathbf{g}}$ is automatically revealed so its revealment is 1. Also,

$$\text{Inf}_{\mathbf{g}}(f) \leq \mathbb{P}_\lambda[\exists(z, R) \in \eta \text{ with } R \geq L \text{ and } \mathbb{B}_R^z \cap \mathbb{B}_r \neq \emptyset] \quad (8)$$

so that this quantity tends to 0 as L tends to infinity (thanks to the moment assumption on μ).

Let us now bound the revealment for $(x, n) \in I_L$. The random variable $\eta_{(x,n)}$ is revealed by $\mathbb{T}_{s,L}$ if the Euclidean distance between \mathbb{S}^x and the connected component of $\partial \mathbb{B}_s$ is smaller than n . Let \mathbb{S}_n^x be the union of the boxes \mathbb{S}^y that contain a point at distance exactly n from \mathbb{S}^x . We deduce that

$$\delta_{(x,n)}(\mathbb{T}_{s,L}) \leq \mathbb{P}_\lambda[\mathbb{S}_n^x \longleftrightarrow \partial \mathbb{B}_s]. \quad (9)$$

Overall, plugging the previous bounds (8) and (9) on the revealments of the algorithms $\mathbb{T}_{s,L}$ into (7), we obtain

$$\theta_r(1 - \theta_r) \leq 2 \sum_{(x,n) \in I_L} \mathbb{P}_\lambda[\mathbb{S}_n^x \longleftrightarrow \partial \mathbb{B}_s] \cdot \text{Inf}_{(x,n)}(f) + o(1).$$

(Above, $o(1)$ denotes a quantity tending to 0 as L tends to infinity.) Letting L tend to infinity (note that the influence $\text{Inf}_{(x,n)}$ does not depend on the algorithms $\mathbb{T}_{s,L}$), we find

$$\theta_r(1 - \theta_r) \leq 2 \sum_{\substack{x \in \mathbb{Z}^d \\ n \geq 1}} \mathbb{P}_\lambda[\mathbb{S}_n^x \longleftrightarrow \partial \mathbb{B}_s] \cdot \text{Inf}_{(x,n)}(f). \quad (10)$$

In order to conclude the proof, we need the following two lemmata. First, a simple union bound argument allows us to bound $\mathbb{P}_\lambda[\mathbb{S}_n^x \longleftrightarrow \partial \mathbb{B}_s]$ in terms of the one-arm probability. More precisely, consider the following lemma, which will be proved at the end of the section.

Lemma 10. *Fix $\lambda_0 > 0$. There exists $c_2 > 0$ such that for every $\lambda \geq \lambda_0$, $x \in \mathbb{Z}^d$ and $n \geq 1$,*

$$\int_0^r \mathbb{P}_\lambda[\mathbb{S}_n^x \longleftrightarrow \partial \mathbb{B}_s] ds \leq c_2 n^{d-1} \Sigma_r(\lambda).$$

Integrating (10) for radii between 0 and r and using the lemma above gives

$$r\theta_r(1 - \theta_r) \leq 2c_2 \Sigma_r \sum_{\substack{x \in \mathbb{Z}^d \\ n \geq 1}} n^{d-1} \text{Inf}_{(x,n)}(f). \quad (11)$$

The most delicate step is to relate the influences in the equation above with the pivotal probabilities appearing in the derivative formula (Russo). This is the content of the following lemma, which will be proved in Section 3.3.

Lemma 11. *There exists $c_3 > 0$ such that for every $x \in \mathbb{Z}^d$, every $n \geq 1$ and every $\lambda > \tilde{\lambda}_c$,*

$$\sum_{x \in \mathbb{Z}^d} \text{Inf}_{(x,n)}(f) \leq c_3 n^{4d-2} \mu[n-1, n] \sum_{x \in \mathbb{Z}^d} \mathbb{E}_\lambda[\text{Piv}_{x,A}].$$

Dividing (11) by Σ_r and applying the lemma above gives

$$\begin{aligned} \frac{r}{\Sigma_r} \theta_r(1 - \theta_r) &\leq 2c_2 c_3 \sum_{n \geq 1} n^{5d-3} \mu[n-1, n] \sum_{x \in \mathbb{Z}^d} \mathbb{E}_\lambda[\text{Piv}_{x,A}] \\ &\leq c_1 \sum_{x \in \mathbb{Z}^d} \mathbb{E}_\lambda[\text{Piv}_{x,A}] \stackrel{(\text{Russo})}{=} c_1 \theta'_r. \end{aligned}$$

This implies Lemma 5. Before proving lemmata 10 and 11, we would like to make several remarks concerning the proof above.

Remark 12. *The interest of working with $\mathbb{T}_{s,L}$ is to have a finite number of coordinates for the algorithm. We could have stated an OSSS inequality valid for countably many states and get (10) directly, but we believe the previous strategy to be shorter and thriftier.*

Remark 13. *Lemma 10 may a priori be improved. Indeed, the union bound in the first inequality is quite wasteful. Nonetheless, with the moment assumption on the radius distribution, the claim above is sufficient.*

Remark 14. Lemma 11 is slightly too strong. We could replace n^{-2d-1} by any sequence a_n such $a_n n^{2d-1}$ is summable. If Lemma 10 was improved by replacing n^{d-1} by a sequence b_n going to infinity more slowly, then the condition on a_n would become that $n^d a_n b_n$ is summable. This is an observation to keep in mind if one wants to improve the moment assumption. Here, we favored the simplest proof possible and did not try to optimize the two lemmata.

Remark 15. We refer to $\lambda \geq \lambda_0$ in Lemma 10 instead of $\lambda > \tilde{\lambda}_c$ (even though the lemma is anyway used in this context) to illustrate that the only place where $\lambda > \tilde{\lambda}_c$ is used is in Lemma 11.

We finish this section with the proof of Lemma 10.

Proof of Lemma 10. Let Y be the subset of \mathbb{Z}^d such that $\mathbf{S}_n^x = \bigcup_{y \in Y} \mathbf{S}^y$. We have that

$$\mathbb{P}_\lambda[\mathbf{S}_n^x \leftrightarrow \partial \mathbf{B}_s] \leq \sum_{y \in Y} \mathbb{P}_\lambda[\mathbf{S}^y \leftrightarrow \partial \mathbf{B}_s] \leq \frac{1}{c_{\text{IT}}} \sum_{y \in Y} \mathbb{P}_\lambda[y \leftrightarrow \partial \mathbf{B}_s].$$

where in the last inequality, we used the FKG inequality, (IT) and the fact that if $\mathbf{S}^y \leftrightarrow \partial \mathbf{B}_s$ and the event \mathcal{D}_y defined below (IT) occur, then y is connected to $\partial \mathbf{B}_s$.

Integrating on s between 0 and r and observing that y is at distance $|s - \|y\||$ of $\partial \mathbf{B}_s$, we deduce that

$$\int_0^r \mathbb{P}_\lambda[y \leftrightarrow \partial \mathbf{B}_s] ds \leq \int_0^r \theta_{|s - \|y\|}(\lambda) ds \leq 2\Sigma_r(\lambda).$$

Since the cardinality of Y is bounded by a constant times n^{d-1} , the result follows. \square

3.2 A technical statement regarding connection probabilities above $\tilde{\lambda}_c$

The following lemma will be instrumental in the proof of Lemma 11. It is the unique place where we use the assumption $\lambda > \tilde{\lambda}_c$.

Below, $X \overset{Z}{\longleftrightarrow} Y$ means that X is connected to Y in $\mathcal{O}(\eta^Z)$, where η^Z is the set of $(z, R) \in \eta$ such that $\mathbf{B}_z(R) \subset Z$. We highlight that this is not the same as the existence of a continuous path in $\mathcal{O}(\eta) \cap Z$ connecting x and y , since such a path could a priori pass through regions in Z which are only covered by balls intersecting $\mathbb{R}^d \setminus Z$.

Lemma 16. *There exists a constant $c_4 > 0$ such that for every $\lambda > \tilde{\lambda}_c$ and $r \geq \mathbf{r}^*$,*

$$\mathbb{P}_\lambda[0 \overset{\mathbf{B}_r}{\longleftrightarrow} \mathbf{B}_{\mathbf{r}^*}^x] \geq \frac{c_4}{r^{2d-2}} \quad \text{for every } x \in \partial \mathbf{B}_r. \quad (12)$$

Remark 17. *Before diving into the proof, let us first explain how we will use the assumption $\lambda > \tilde{\lambda}_c$. Fix $r > 0$. If \mathbf{B}_r is connected to $\partial \mathbf{B}_{2r}$, then there must exist $x \in \mathbb{Z}^d$ with $\mathbf{S}^x \cap \partial \mathbf{B}_r \neq \emptyset$ such that \mathbf{S}^x is connected to $\partial \mathbf{B}_{2r}$. Since above $\tilde{\lambda}_c$, the probability of the former is bounded away from 0 uniformly in r , and since the probabilities of the latter events are all smaller than $\mathbb{P}_\lambda[\mathbf{S}^0 \leftrightarrow \partial \mathbf{B}_r]$, the union bound gives*

$$\mathbb{P}_\lambda[\mathbf{S}^0 \leftrightarrow \partial \mathbf{B}_r] \geq \frac{c_5}{r^{d-1}}. \quad (13)$$

The argument will rely on the following observation. As explained in (13), the probability that S^0 is connected to ∂B_r does not decay quickly. This event implies the existence of y such that S^0 is connected in B_r to $B_{r_*}^y$, and one ball B_R^z (with $(z, R) \in \eta$) covering $B_{r_*}^y$ and intersecting the complement of B_r . The problem is that this site y may be quite far from ∂B_r . Nonetheless, this seems unlikely since the cost (by the moment assumption on μ) of having a large ball B_R^z intersecting $B_{r_*}^y$ is overwhelmed by the probability that the latter is connected to ∂B_r in $B_{r-\|y\|}^y$ by a path in $\mathcal{O}(\eta)$ using a priori smaller balls (and maybe even only balls included in B_r). The proof below will harvest this idea though with some important variations due to the fact that all the balls under consideration must remain inside B_r . One of the key idea is the introduction of a new scale \mathbf{r}^{**} and an induction on the probability that S^0 is connected to $B_{\mathbf{r}^{**}}^x$ in B_r , where $x \in \partial B_r$.

Proof. We will prove that there exist $\mathbf{r}^{**} > 0$ and $c_6 > 0$ such that

$$\mathbb{P}_\lambda[S^0 \xleftrightarrow{B_r} B_{\mathbf{r}^{**}}^x] \geq \frac{c_6}{r^{2d-2}} \quad \text{for every } x \in \partial B_r. \quad (14)$$

This is amply sufficient to prove the lemma since, if T denotes the set of $t \in \mathbb{Z}^d$ such that $S^t \cap B_{2\mathbf{r}^{**}}^x \neq \emptyset$ and $B_{\mathbf{r}^{**}}^t \subset B_r$, then

$$\mathbb{P}_\lambda[0 \xleftrightarrow{B_r} B_{\mathbf{r}^{**}}^x] \stackrel{\text{(FKG)}}{\geq} \mathbb{P}_\lambda[\mathcal{D}_0] \left(\prod_{t \in T} \mathbb{P}_\lambda[\mathcal{D}_t] \right) \mathbb{P}_\lambda[S^0 \xleftrightarrow{B_r} B_{\mathbf{r}^{**}}^x] \stackrel{\text{(IT)}}{\geq} c_{\text{IT}}^{|T|+1} \cdot \frac{c_6}{r^{2d-2}}.$$

We therefore focus on the proof of (14). We will fix $\mathbf{r}^{**} \geq 2\mathbf{r}^*$ sufficiently large (see the end of the proof). Introduce $Y := \mathbb{Z}^d \cap B_{r-\mathbf{r}^{**}}$ and a finite set $Z \subset \partial B_r$ such that the union of the balls $B_{r_*}^y$ and $B_{\mathbf{r}^{**}}^z$ with $y \in Y$ and $z \in Z$ cover the ball B_r . Note that if 0 is connected to ∂B_r , then either one of the $z \in Z$ is such that 0 is connected to $B_{\mathbf{r}^{**}}^z$ in B_r , or there exists $y \in Y$ such that the event

$$\mathcal{A}(y) := \{S^0 \xleftrightarrow{B_r} B_{r_*}^y\} \cap \{\exists (u, R) \in \eta \text{ such that } B_R^u \text{ intersects both } B_{r_*}^y \text{ and } \partial B_r\}$$

occurs. The union bound therefore implies that

$$\sum_{z \in Z} \mathbb{P}_\lambda[S^0 \xleftrightarrow{B_r} B_{\mathbf{r}^{**}}^z] + \sum_{y \in Y} \mathbb{P}_\lambda[\mathcal{A}(y)] \geq \mathbb{P}_\lambda[S^0 \xleftrightarrow{\partial B_r}] \stackrel{(13)}{\geq} \frac{c_5}{r^{d-1}}. \quad (15)$$

For $y \in Y$, introduce $x := (r/\|y\|)y \in \partial B_r$ and $\bar{r} = r - \|y\|$. The event on the right of the definition of $\mathcal{A}(y)$ is independent of the event on the left. Since any point in $B_{r_*}^y$ is at a distance at least $\bar{r} - \mathbf{r}_*$ of ∂B_r , we deduce that

$$\mathbb{P}_\lambda[\mathcal{A}(y)] \leq \left(c_7 \int_{\bar{r}-\mathbf{r}_*}^{\infty} r^{d-1} \mu(dr) \right) \cdot \mathbb{P}_\lambda[S^0 \xleftrightarrow{B_r} B_{r_*}^y] \leq \frac{c_8}{\bar{r}^{4d-2}} \mathbb{P}_\lambda[S^0 \xleftrightarrow{B_r} B_{r_*}^y].$$

In the second inequality, we used the moment assumption on μ and the fact that $\bar{r} - \mathbf{r}_* \geq \bar{r}/2$ since $\mathbf{r}^{**} \geq 2\mathbf{r}^*$. Using this latter assumption one more time implies that

$$\begin{aligned} \mathbb{P}_\lambda[S^0 \xleftrightarrow{B_r} B_{\mathbf{r}^{**}}^x] &\stackrel{\text{(FKG)}}{\geq} \mathbb{P}_\lambda[S^0 \xleftrightarrow{B_r} B_{r_*}^y] \cdot \mathbb{P}_\lambda[\mathcal{D}_y] \cdot \mathbb{P}_\lambda[S^y \xleftrightarrow{B_{\bar{r}}} B_{\mathbf{r}^{**}}^x] \\ &\geq \frac{c_{\text{IT}} \bar{r}^{4d-2}}{c_8} \cdot \mathbb{P}_\lambda[\mathcal{A}(y)] \cdot \mathbb{P}_\lambda[S^y \xleftrightarrow{B_{\bar{r}}} B_{\mathbf{r}^{**}}^x]. \end{aligned} \quad (16)$$

From now on, define $U(r) := \mathbb{P}_\lambda[S^0 \xleftrightarrow{B_r} B_{\mathbf{r}^{**}}^x]$ for $x \in \partial B_r$ (the choice of x is irrelevant by invariance under the rotations). Now, one may choose Z in such a way that $|Z| \leq c_9 r^{d-1}$. Plugging (16) in (15), we obtain the following inequality

$$c_9 r^{d-1} U(r) + c_{10} \sum_{y \in Y} \frac{U(r)}{U(r - \|y\|)(r - \|y\|)^{4d-2}} \geq \frac{c_5}{r^{d-1}}.$$

Using that $U(r) = 1$ for every $r \leq \mathbf{r}^{**}$, we deduce (14) by induction, provided \mathbf{r}^{**} is chosen large enough to start with. \square

3.3 Proof of Lemma 11

We are now in a position to prove Lemma 11. Fix $x \in \mathbb{Z}^d$ and $r, n \geq 1$. Define

$$\mathcal{P}_x(n) := \{0 \longleftrightarrow B_n^x\} \cap \{B_n^x \longleftrightarrow \partial B_r\} \cap \{0 \longleftrightarrow \partial B_r\}^c.$$

Introduce $\mathbf{C} = \mathbf{C}(\eta) := \{z \in \mathbb{R}^d : z \longleftrightarrow 0\}$ and $\mathbf{C}' = \mathbf{C}'(\eta) := \{z \in \mathbb{R}^d : z \longleftrightarrow \partial B_r\}$ the connected components of 0 and ∂B_r in $\mathcal{O}(\eta)$. Our first goal is to prove that, conditionally on $\mathcal{P}_x(n)$, the probability that \mathbf{C} and \mathbf{C}' are close to each other in B_{3n}^x is not too small.

Claim. *We have that*

$$\mathbb{P}_\lambda[\mathcal{P}_x(n) \text{ and } d(\mathbf{C} \cap B_{3n}^x, \mathbf{C}') < \mathbf{r}^*] \geq \frac{c_{11}}{n^{3d-2}} \cdot \mathbb{P}_\lambda[\mathcal{P}_x(n)].$$

Before proving this claim, let us conclude the proof of the lemma. If the event on the left occurs, there must exist $y \in \mathbb{Z}^d$ within a distance at most \mathbf{r}_* of both $\mathbf{C} \cap B_{3n}^x$ and \mathbf{C}' : simply pick a point $z \in \mathbb{R}^d$ at a distance smaller than $\mathbf{r}^*/2$ of both $\mathbf{C} \cap B_{3n}^x$ and \mathbf{C}' , and then take $y \in \mathbb{Z}^d$ within distance \sqrt{d} of it (we used the inequality on the right in (5)). In particular, $\mathcal{P}_y(\mathbf{r}_*)$ must occur for this y and we deduce that

$$\mathbb{P}_\lambda[\mathcal{P}_x(n)] \leq c_{12} n^{3d-2} \sum_{\substack{y \in \mathbb{Z}^d \\ \|y-x\| \leq 3n+\mathbf{r}_*}} \mathbb{P}_\lambda[\mathcal{P}_y(\mathbf{r}_*)]. \quad (17)$$

To conclude, observe that the definition of \mathbf{r}^* given by (IT) implies that for all y ,

$$c_{\text{IT}} \mathbb{P}_\lambda[\mathcal{P}_y(\mathbf{r}_*)] \leq \mathbb{E}_\lambda[\text{Piv}_{y,A}]. \quad (18)$$

Now, one easily gets that

$$\text{Inf}_{(x,n)}(f) \leq \lambda \mu[n-1, n] \times \mathbb{P}_\lambda[\mathcal{P}_x(n + \sqrt{d})]. \quad (19)$$

(Simply observe that $\mathcal{P}_x(n + \sqrt{d})$ must occur in η , and that the resampled Poisson point process $\tilde{\eta}_{(x,n)}$ must contain at least one point). Plugging (17) (applied to $n + \sqrt{d}$) in (19), then using (18), and finally summing over every $x \in \mathbb{Z}^d$ gives the lemma.

Proof of the claim. The proof consists in expressing the probability that \mathbf{C} and \mathbf{C}' come within a distance \mathbf{r}^* of each other in terms of the probability that they remain at a distance at least \mathbf{r}^* of each other.

Fix $y \in \mathbb{Z}^d$. For a subset C of \mathbb{R}^d , let $u_C = u_C(y)$ be the point of S^y furthest to C , and $v_C = v_C(y)$ the point of C closest to u_C . Consider the event

$$\mathcal{E}_y := \{\mathbf{C} \cap \mathbb{B}_n^x \neq \emptyset \text{ and } d(u_{\mathbf{C}}, \mathbf{C}) \geq \mathbf{r}^*\}.$$

Note that the event \mathcal{E}_y is measurable in terms of \mathbf{C} . Let us study, on \mathcal{E}_y , the conditional expectation with respect to \mathbf{C} . Introduce the three events

$$\mathcal{F}_y := \mathcal{D}_{u_{\mathbf{C}}} \quad \mathcal{G}_y := \{u_{\mathbf{C}} \overset{\mathbb{R}^d \setminus \mathbf{C}}{\longleftrightarrow} \mathbb{B}_{v_{\mathbf{C}}}(\mathbf{r}^*)\} \quad \mathcal{H}_y := \{\mathbb{B}_n^x \cap S^y \overset{\mathbb{R}^d \setminus \mathbf{C}}{\longleftrightarrow} \partial \mathbb{B}_r\}.$$

On the event \mathcal{E}_y , conditioned on \mathbf{C} , $\eta^{\mathbb{R}^d \setminus \mathbf{C}}$ has the same law as $\tilde{\eta}^{\mathbb{R}^d \setminus \mathbf{C}}$ for some independent realization $\tilde{\eta}$ of the Poisson point process (recall the definition of η^Z from the previous section). Also, the distance between $u_{\mathbf{C}}$ and $v_{\mathbf{C}}$ (or equivalently \mathbf{C}) is larger than \mathbf{r}^* . Therefore, we may use (IT), Lemma 16 and the FKG inequality to get the following inequality between conditional expectations:

$$\mathbb{P}_\lambda[\mathcal{F}_y \cap \mathcal{G}_y \cap \mathcal{H}_y | \mathbf{C}] \geq \mathbb{P}_\lambda[\mathcal{F}_y | \mathbf{C}] \cdot \mathbb{P}_\lambda[\mathcal{G}_y | \mathbf{C}] \cdot \mathbb{P}_\lambda[\mathcal{H}_y | \mathbf{C}] \geq c_{\text{IT}} \frac{c_4}{n^{2d-2}} \mathbb{P}_\lambda[\mathcal{H}_y | \mathbf{C}] \quad \text{a.s. on } \mathcal{E}_y. \quad (20)$$

Now, observe that if \mathcal{E}_y and \mathcal{H}_y occur simultaneously, then $\mathcal{P}_x(n)$ occurs (we use $\mathbf{r}^* \geq \sqrt{d}$). Furthermore if \mathcal{F}_y and \mathcal{G}_y also occur simultaneously with them, then $\mathbb{B}_{\mathbf{r}^*}^{v_{\mathbf{C}}} \cap \mathbf{C}' \neq \emptyset$. Since by construction $v_{\mathbf{C}} \in \mathbb{B}_{3n}^x$, we deduce that $d(\mathbf{C} \cap \mathbb{B}_{3n}^x, \mathbf{C}') < \mathbf{r}^*$. Integrating (20) on \mathcal{E}_y , we deduce that

$$\mathbb{P}_\lambda[\mathcal{P}_x(n) \text{ and } d(\mathbf{C} \cap \mathbb{B}_{3n}^x, \mathbf{C}') < \mathbf{r}^*] \geq \mathbb{P}_\lambda[\mathcal{E}_y \cap \mathcal{F}_y \cap \mathcal{G}_y \cap \mathcal{H}_y] \geq \frac{c_{\text{IT}} c_4}{n^{2d-2}} \mathbb{P}_\lambda[\mathcal{E}_y \cap \mathcal{H}_y].$$

Observe that if $\mathcal{P}_x(n)$ and $d(\mathbf{C} \cap \mathbb{B}_n^x, \mathbf{C}') \geq \mathbf{r}^*$ occur, then there exists $y \in \mathbb{Z}^d$ such that the event on the right-hand side occurs. In particular, S^y must intersect \mathbb{B}_n^x for \mathcal{H}_y to occur, so that there are $c_6 n^d$ possible values for y . Summing over all these values, we therefore get that

$$c_6 n^d \mathbb{P}_\lambda[\mathcal{P}_x(n) \text{ and } d(\mathbf{C} \cap \mathbb{B}_{3n}^x, \mathbf{C}') < \mathbf{r}^*] \geq \frac{c_{\text{IT}} c_4}{n^{2d-2}} \mathbb{P}_\lambda[\mathcal{P}_x(n) \text{ and } d(\mathbf{C} \cap \mathbb{B}_n^x, \mathbf{C}') \geq \mathbf{r}^*],$$

which implies the claim readily. \square

4 Renormalization of crossing probabilities

In this section, we prove Theorems 2 and 3. The proof is quite different in the regime where $\mu[r, \infty]$ decays exponentially fast (light tail), and in the regime where it does not (heavy tail). Also, since in the heavy tail regime the renormalization argument performed

below may be of some use in the study of other models, or for different distributions μ , we prove a quantitative lemma which we believe to be of independent interest.

In this section, we reboot the count for the constants c_i starting from c_1 . The section is organized as follows. In Section 4.1, we prove a renormalization lemma which will enable us to derive the theorems. In Sections 4.2 and 4.3, we derive Theorem 2 and Theorem 3 respectively.

4.1 The renormalization lemma

Introduce, for every $\delta, \alpha, \lambda \geq 0$ and $r \geq 1$, the two functions

$$\begin{aligned}\pi_r^\delta(\lambda) &:= \mathbb{P}_\lambda[\exists(z, R) \in \eta \text{ such that } \mathbf{B}_R^z \cap \mathbf{B}_{2\delta r} \neq \emptyset \text{ and } \mathbf{B}_R^z \cap \partial\mathbf{B}_{(1-2\delta)r} \neq \emptyset], \\ \theta_r^\alpha(\lambda) &:= \mathbb{P}_\lambda[\mathbf{B}_{\alpha r} \longleftrightarrow \partial\mathbf{B}_r].\end{aligned}$$

Note that $\pi_r^0(\lambda) = \phi_r(\lambda)$ and $\theta_r^0(\lambda) = \theta_r(\lambda)$ by definition.

Remark 18. *The quantity $\pi_r^\delta(\lambda)$ is expressed in terms of μ as follows: let c_d be the area of the sphere of radius 1 in \mathbb{R}^d , then*

$$\pi_r^\delta(\lambda) = 1 - \exp\left(-\lambda c_d \int_0^\infty a^{d-1} \mu[|a - 2\delta r| \vee |r - 2\delta r - a|, \infty] da\right). \quad (21)$$

The following lemma will be the key to the proofs of the theorems.

Lemma 19 (Renormalization inequality). *For every $0 < \alpha \leq \delta \leq 1/4$, there exists $c_1 > 0$ such that for every $\lambda, r > 0$,*

$$\theta_r^\alpha(\lambda) \leq \pi_r^\delta(\lambda) + \frac{c_1}{(\delta^2 \alpha)^d} \max_{\substack{u, v \geq \delta \\ u+v=1-\alpha}} \theta_{ur}^\alpha(\lambda) \theta_{vr}^\alpha(\lambda). \quad (22)$$

Note that the smaller the α and δ , the larger the entropic factor $c_1/(\delta^2 \alpha)^d$. We will see that the choices of α and δ are important in the applications. Except for the exponential bound on $\theta_r(\lambda)$ in the proof of Theorem 2, we will always pick $\delta = \alpha$.

Proof. Fix $0 < \alpha \leq \delta \leq 1/4$ and $\lambda, r > 0$. Set

$$E := \{2\delta r\} \cup ([2\delta r, (1-2\delta)r] \cap \frac{\alpha\delta}{2} r\mathbb{Z}) \cup \{(1-2\delta)r\}$$

and index the elements of E in the increasing order by $r_0 < r_1 < \dots < r_\ell$. Recall the notation η^Z for the set of $(z, R) \in \eta$ with $\mathbf{B}_R^z \subset Z$, and introduce the three events

$$\begin{aligned}\mathcal{A} &:= \{\exists(z, R) \in \eta : \mathbf{B}_R^z \cap \mathbf{B}_{2\delta r} \neq \emptyset \text{ and } \mathbf{B}_R^z \cap \partial\mathbf{B}_{(1-2\delta)r} \neq \emptyset\}, \\ \mathcal{B}_k &:= \{\mathbf{B}_{\alpha r} \longleftrightarrow \partial\mathbf{B}_{r_k} \text{ in } \mathcal{O}(\eta^{\mathbf{B}_{r_{k+1}}})\}, \\ \mathcal{C}_k &:= \{\mathbf{B}_{r_{k+1}} \longleftrightarrow \partial\mathbf{B}_r\}.\end{aligned}$$

If $\mathbf{B}_{\alpha r}$ is connected to $\partial\mathbf{B}_r$ but \mathcal{A} does not occur, then $\mathcal{B}_k \cap \mathcal{C}_k$ must occur for some $0 \leq k < \ell$ (we use that $\delta \geq \alpha$). Furthermore, by construction, \mathcal{B}_k and \mathcal{C}_k are independent

since \mathcal{C}_k is measurable with respect to $\mathcal{O}(\eta \setminus \eta^{B_{r_k}})$. These two observations together imply that

$$\theta_r^\alpha(\lambda) \leq \mathbb{P}_\lambda[\mathcal{A}] + \sum_{k=0}^{\ell-1} \mathbb{P}_\lambda[\mathcal{B}_k \cap \mathcal{C}_k] = \pi_r^\delta(\lambda) + \sum_{k=0}^{\ell-1} \mathbb{P}_\lambda[\mathcal{B}_k] \mathbb{P}_\lambda[\mathcal{C}_k]. \quad (23)$$

Fix $0 \leq k < \ell$. Let X denote a set of points $x \in \partial B_{\alpha r}$ such that the union of the balls $B_{\alpha \delta r}^x$ for $x \in X$ covers $\partial B_{\alpha r}$. Choose X such that $|X| \leq c_2 \delta^{-(d-1)}$. Applying the union bound, we find

$$\mathbb{P}_\lambda[\mathcal{B}_k] \leq \sum_{x \in X} \mathbb{P}_\lambda[B_{\alpha \delta r}^x \leftrightarrow \partial B_{r_k - \alpha r}^x] \leq |X| \theta_{r_k - \alpha r}^\alpha(\lambda) \leq \frac{c_2}{\delta^{d-1}} \theta_{r_k - \alpha r}^\alpha(\lambda). \quad (24)$$

Similarly, using a covering of $\partial B_{r_{k+1}}$ with at most $c'_2(\alpha \delta)^{1-d}$ balls of radius $\alpha \delta r$ centered on ∂B_{r_k} , we obtain

$$\mathbb{P}_\lambda[\mathcal{C}_k] \leq \frac{c'_2}{(\delta \alpha)^{d-1}} \theta_{r-r_k}^\alpha(\lambda). \quad (25)$$

Plugging (24) and (25) in (23) and using the fact that $\ell \leq 2/(\alpha \delta)$ concludes the proof. \square

The previous lemma leads to the following useful corollary.

Corollary 20. For $\lambda < \tilde{\lambda}_c$, $\lim_{r \rightarrow \infty} \mathbb{P}_\lambda[B_r \leftrightarrow \partial B_{2r}] = 0$.

Proof. Choose $\alpha = 1/6$. In this case

$$\theta_r^\alpha(\lambda) = \mathbb{P}_\lambda[B_{r/6} \leftrightarrow \partial B_r].$$

Using a covering of ∂B_r with balls of radius $B_{r/6}$, it suffices to prove that $\theta_r^\alpha(\lambda)$ converges to 0. A reasoning similar to the bound on $\mathbb{P}_\lambda[\mathcal{B}_k]$ in the last proof implies that for every $s \in [\alpha r, r]$,

$$\theta_r^\alpha(\lambda) \leq c_3 \theta_s^\alpha(\lambda), \quad (26)$$

which enables us to rewrite (22) (applied with $\delta = \alpha = 1/6$) as

$$\theta_r^\alpha(\lambda) \leq \pi_r^\alpha(\lambda) + c_4 \theta_{\alpha r}^\alpha(\lambda)^2.$$

Now, fix $\varepsilon > 0$ such that $2c_4\varepsilon < 1$ and work with r_0 large enough that $\pi_r^\alpha(\lambda) \leq \varepsilon/2$ for every $r \geq r_0$. Since $\lambda < \tilde{\lambda}_c$, we can pick $r \geq r_0$ such that $\theta_r^\alpha(\lambda) < \varepsilon$. We deduce inductively that

$$\theta_{r/\alpha^k}^\alpha(\lambda) \leq \varepsilon$$

for every $k \geq 1$. Using (26) one last time gives that $\theta_s^\alpha(\lambda) \leq c_3\varepsilon$ for every $s \geq r$. \square

4.2 Proof of Theorem 2

Consider $\lambda < \tilde{\lambda}_c$. Fix $\delta \in (0, \frac{1}{2})$ and observe that since $\mu[r, \infty] \leq \exp(-cr)$, (21) implies that there exists $c' > 0$ such that $\pi_r^\delta(\lambda) \leq \exp(-c'r)$ for every $r \geq 1$. We now proceed in two steps: we first show that $\theta_r(\lambda) \leq r^{-d-1}$ for r large enough, and then improve this estimate to an exponential decay.

Polynomial bound on $\theta_r(\lambda)$ Lemma 19 applied to $\alpha = \delta = 1/6$ implies that for every $r \geq 1$

$$\theta_r^\alpha(\lambda) \leq e^{-c'r} + c_5 \max_{\substack{u, v \geq \alpha \\ u+v=1-\alpha}} \theta_{ur}^\alpha(\lambda) \theta_{vr}^\alpha(\lambda). \quad (27)$$

Since $\lambda < \tilde{\lambda}_c$, Corollary 20 implies that $\theta_r^\alpha(\lambda)$ converges to 0 as r tends to infinity. In particular, we deduce that for every $\varepsilon > 0$,

$$\theta_r^\alpha(\lambda) \leq e^{-c'r} + \varepsilon \theta_{\alpha r}^\alpha(\lambda)$$

for r large enough. A simple induction implies that $\theta_r^\alpha(\lambda) \leq r^{-d-1}$ for r large enough.

Exponential bound on $\theta_r(\lambda)$ Choose $\delta = 1/6$. For each $r \geq 6$, apply Lemma 19 with $\alpha = \alpha(r) := 1/r$ to get that

$$\theta_r(\lambda) \leq e^{-c'r} + c_6 r^d \max_{\substack{u, v \geq \frac{1}{6} \\ u+v=1-\alpha(r)}} \theta_{ur}(\lambda) \theta_{vr}(\lambda),$$

where we use that

$$\theta_r(\lambda) \leq \theta_r^{\alpha(r)}(\lambda) = \mathbb{P}_\lambda[\mathbf{B}_1 \longleftrightarrow \partial \mathbf{B}_r] \leq \frac{1}{c_{\text{IT}}} \theta_r(\lambda).$$

Set $c_7 := 2c_6 e$ and consider r_0 large enough so that for every $r \geq \delta r_0$,

$$\begin{aligned} c_7 r^d e^{-c'r} &\leq \frac{1}{2} e^{-r/r_0}, \text{ and} \\ c_7 r^d \theta_r(\lambda) &\leq e^{-1}. \end{aligned}$$

(The second constraint is satisfied thanks to the polynomial bound derived in the first part of the proof.) By induction on k , one can show that for every $k \geq 0$,

$$\forall r \in [\delta r_0, r_0/(1-\delta)^k] \quad c_7 r^d \theta_r(\lambda) \leq \exp(-r/r_0).$$

4.3 Proof of Theorem 3

In this section, we prove Theorem 3 by proving the following quantitative lemma. We believe the lemma to be of value for distributions μ other than the one considered in Theorem 3. Before that, note that the lower bound

$$\theta_r(\lambda) \geq \phi_r(\lambda) \quad (28)$$

for every $r > 0$ is trivial since 0 is connected to $\partial \mathbf{B}_r$ in the case where there exists $(z, R) \in \eta$ such that \mathbf{B}_R^z contains 0 and intersects $\partial \mathbf{B}_r$. Therefore, the main concern of this section will be an upper bound on $\theta_r^\alpha(\lambda)$ (which is larger than $\theta_r(\lambda)$).

Lemma 21. Consider $\lambda > 0$, $\eta < 1$ and $\alpha \in (0, \frac{1}{4})$ small enough such that $\alpha^\eta + (1-\alpha)^\eta \geq 1$. For every $\varepsilon > 0$ sufficiently small and $r \geq 1$, if there exists $r_0 \leq \alpha r$ such that

$$\theta_s^\alpha(\lambda) \leq \varepsilon \pi_r^\alpha(\lambda)^{(s/r)^\eta} \quad \forall s \in [r_0, r_0/\alpha] \quad (29)$$

$$\pi_s^\alpha(\lambda) \leq \varepsilon \pi_r^\alpha(\lambda)^{(s/r)^\eta} \quad \forall s \in [r_0, (1-\alpha)r], \quad (30)$$

then

$$\theta_r^\alpha(\lambda) \leq (1 + \varepsilon) \pi_r^\alpha(\lambda). \quad (31)$$

Remark 22. 1) Note that distributions μ with exponential decay do not satisfy the assumptions of the lemma for any r , provided that ε is small enough.

2) For distributions μ satisfying $\lim_{r \rightarrow \infty} \pi_r^\alpha(\lambda)^{1/r^\eta} = \kappa$, for some constant κ , the existence of r_0 satisfying (29) when $\lambda < \tilde{\lambda}_c$ follows directly from

- picking r_0 such that $\theta_r^\alpha(\lambda) < \frac{\varepsilon}{2\kappa}$ for every $r \geq r_0$ (by Corollary 20),
- picking r large enough that $\pi_r^\alpha(\lambda)^{1/r^\eta} \geq \kappa 2^{-(\alpha/r_0)^\eta}$.

3) The second assumption (30) is a regularity statement on $\pi_r^\alpha(\lambda)$ that can be obtained from the regularity of $\mu[r, \infty]$ using (21).

The proof of the lemma will be based on the recursive relation given by Lemma 19. We use a strategy which is inspired by the study of differential equations. We control the function $f(s) := \theta_s^\alpha(\lambda)$ in terms of $g(s) := \pi_r^\alpha(\lambda)^{(s/r)^\eta}$ uniformly for $s \in [r_0, (1-\alpha)r]$ to finally deduce a bound for $f(r)$ using again Lemma 19.

Proof. Set $C = c_1 \alpha^{-2d}$, where c_1 is given by Lemma 19, and assume that $\varepsilon < \frac{1}{4C}$. The key ingredient will be the following inequality: for every $s \in [r_0, (1-\alpha)r]$,

$$\theta_s^\alpha(\lambda) \leq 2\varepsilon \pi_r^\alpha(\lambda)^{(s/r)^\eta}. \quad (32)$$

For $s \in [r_0, r_0/\alpha]$, (32) follows from (29). To obtain the claim for $s > r_0/\alpha$, use the induction hypothesis in the second line below to get

$$\begin{aligned} \theta_s^\alpha(\lambda) &\stackrel{(22)}{\leq} \pi_s^\alpha(\lambda) + C \max_{\substack{u, v \geq \alpha \\ u+v=1-\alpha}} \theta_{us}^\alpha(\lambda) \theta_{vs}^\alpha(\lambda) \\ &\stackrel{(32)}{\leq} \pi_s^\alpha(\lambda) + 4C\varepsilon^2 \max_{\substack{u, v \geq \alpha \\ u+v=1-\alpha}} \pi_r^\alpha(\lambda)^{(u^\eta+v^\eta)(s/r)^\eta} \\ &\stackrel{(30)}{\leq} (\varepsilon + 4C\varepsilon^2) \pi_r^\alpha(\lambda)^{(s/r)^\eta} \\ &\leq 2\varepsilon \pi_r^\alpha(\lambda)^{(s/r)^\eta}. \end{aligned}$$

In the first line, we applied (22) with $\delta = \alpha$. In the second line, we used (32) for $us, vs \in [r_0, s - r_0]$ and in the third line we used $u^\eta + v^\eta \geq 1$ for every $u, v \geq \alpha$ satisfying

$u + v = 1 - \alpha$. One may apply the first two lines in the previous sequence of inequalities for $s = r$ (since in such case $ur, vr \in [r_0, (1 - \alpha)r]$) so that

$$\theta_r^\alpha(\lambda) \leq \pi_r^\alpha(\lambda) + 4C\varepsilon^2 \max_{\substack{u, v \geq \alpha \\ u+v=1-\alpha}} \pi_r^\alpha(\lambda)^{u^\eta+v^\eta} \leq (1 + \varepsilon) \pi_r^\alpha(\lambda).$$

This concludes the proof. \square

Proof of Theorem 3. Fix $\lambda < \tilde{\lambda}_c$. Since (28) gives the lower bound, we focus on the upper bound. Under the condition C1, define $f(\alpha, r)$ by

$$\pi_r^\alpha(\lambda) = f(\alpha, r)r^{-c}, \quad (33)$$

and, under the condition C2, define $g(\alpha, r)$ by

$$\pi_r^\alpha(\lambda) = g(\alpha, r)r^{d-\gamma} \exp[-c(\frac{r}{2} - 2\alpha r)^\gamma]. \quad (34)$$

Using (21), one can check that $f(\alpha, r)$ and $g(\alpha, r)$ converge (as r tends to infinity) uniformly in $0 \leq \alpha < \frac{1}{4}$ to two continuous positive functions $f(\alpha)$ and $g(\alpha)$.

Consider ε very small and use Corollary 20 to guarantee that $\theta_s^\alpha(\lambda) < \varepsilon$ for every s large enough. Then, for every distribution μ satisfying C1 or C2, one can find $0 < \eta < 1$, r_0 large enough such that (29) and (30) are satisfied for every $r \geq r_0$ large enough, and α small enough (use (33) and (34) and Remark 22 to check the two conditions), so that Lemma 21 implies that for r large enough,

$$\theta_r^\alpha(\lambda) \leq (1 + \varepsilon)\pi_r^\alpha(\lambda). \quad (35)$$

Thanks to (33), we obtain the claim in case C1 by letting ε and then α tend to 0. For case C2, we need to do more, since letting α tend to 0 slowly gives only that $\theta_r(\lambda) \leq \phi_r(\lambda)^{1+o(1)}$. More precisely, we prove the following claim.

Claim *Assume that μ satisfies C2. Fix $\bar{\gamma} \in (\gamma, 1)$. There exist $\alpha > 0$ and $R < \infty$ such that for every $k \geq 0$ and every $r \geq 3^k R$, we have that*

$$\theta_r^{\alpha_k}(\lambda) \leq (1 + 2^{-k})\pi_r^{\alpha_k}(\lambda) \quad (36)$$

where $\alpha_k := \alpha 3^{-k\bar{\gamma}}$.

Before proving the claim, note that it implies, together with (34), that for every $k \geq 0$ and $r \in [3^k R, 3^{k+1} R)$,

$$\theta_r(\lambda) \leq \theta_r^{\alpha_k}(\lambda) \leq (1 + 2^{-k})\pi_r^{\alpha_k}(\lambda) \leq (1 + 2^{-k}) \frac{g(\alpha_k, r)}{g(0, r)} e^{Cr\gamma - \bar{\gamma}} \phi_r(\lambda)$$

for some constant $C = C(\alpha, R) > 0$. This proves the theorem in case C2 thanks to the choice of $\bar{\gamma} > \gamma$ and the uniform convergence of $g(\cdot, r)$ towards a continuous function.

Proof of the Claim. We prove (36) recursively. Fix $0 < \alpha \leq 1/8$ such that

$$2(1 - 12\alpha)^\gamma \left(\frac{1-\alpha}{2}\right)^\gamma \geq 1 + \frac{1}{4}\alpha^\gamma, \text{ and} \quad (37)$$

$$\forall x \in [0, \alpha] \quad (1 - 12x)^{2\gamma} + (x/2)^\gamma \geq 1 + \frac{1}{4}x^\gamma. \quad (38)$$

Fix also $R \geq 1$. There will be several conditions on R that will be added during the proof. The important feature is that every time a new condition is added, one only needs to take R possibly larger than before.

Define M such that for every $x \leq \frac{1}{4}$, $\frac{2}{M} \leq g(x) \leq \frac{M}{2}$. By uniform convergence, R can be chosen large enough that for every $x \leq \frac{1}{4}$ and $r \geq R$,

$$\frac{1}{M} \leq g(x, r) \leq M. \quad (39)$$

Use (35) to choose R in such a way that $\theta_r^\alpha(\lambda) \leq 2\pi_r^\alpha(\lambda)$ for every $r \geq \alpha R$. Then, (36) follows for $k = 0$ by definition. We now assume that (36) is valid for $k - 1$ and prove it for k . Lemma 19 applied to $r \geq R2^k$ and α_k gives that

$$\theta_r^{\alpha_k}(\lambda) \leq \pi_r^{\alpha_k}(\lambda) + \frac{c_1}{\alpha_k^{3d}} \theta_{ur}^{\alpha_k}(\lambda) \theta_{vr}^{\alpha_k}(\lambda), \quad (40)$$

for some $u, v \geq \alpha_k$ such that $u + v = 1 - \alpha_k$. Without loss of generality, we can assume that $u \geq \frac{1}{3}$ and $v \leq \frac{1 - \alpha_k}{2}$. In particular, we have $ur \geq R3^{k-1}$. Also, $vr \geq \alpha R3^{k(1-\gamma)} \geq \alpha R$. The induction hypothesis (in fact we simply bound $2^{-(k-1)}$ by 1) implies that

$$\theta_{ur}^{\alpha_k}(\lambda) \leq \theta_{ur}^{\alpha_{k-1}}(\lambda) \leq 2\pi_{ur}^{\alpha_{k-1}}(\lambda) \quad \text{and} \quad \theta_{vr}^{\alpha_k}(\lambda) \leq \theta_{vr}^\alpha(\lambda) \leq 2\pi_{vr}^\alpha(\lambda).$$

We may use the induction hypothesis to get that

$$\begin{aligned} \theta_{ur}^{\alpha_k}(\lambda) \theta_{vr}^{\alpha_k}(\lambda) &\stackrel{(36)}{\leq} 4\pi_{ur}^{\alpha_{k-1}}(\lambda) \pi_{vr}^\alpha(\lambda) \\ &\stackrel{(39)}{\leq} 4M^2 (uvr^2)^{d-\gamma} \exp\left[-c(r/2)^\gamma \left((u - 4\alpha_{k-1}u)^\gamma + (v - 4\alpha v)^\gamma\right)\right]. \end{aligned} \quad (41)$$

We now bound the term

$$h(v) := (u - 4\alpha_{k-1}u)^\gamma + (v - 4\alpha v)^\gamma = (1 - 4 \cdot 3^{\bar{\gamma}} \alpha_k)^\gamma (1 - \alpha_k - v)^\gamma + (1 - 4\alpha)^\gamma v^\gamma$$

appearing in the exponential in (41). Since $\alpha_k \leq v \leq \frac{1 - \alpha_k}{2}$, an elementary analysis of the function h shows that $h(v) \geq \min(h(\alpha_k), h(\frac{1 - \alpha_k}{2}))$. Using (37) and (38) to bound $h(\frac{1 - \alpha_k}{2})$ and $h(\alpha_k)$ respectively, we obtain

$$h(v) \geq 1 + \frac{1}{4}\alpha_k^\gamma \geq (1 - 4\alpha_k)^\gamma + \frac{1}{4}\alpha_k^\gamma.$$

Plugging this estimate in (41) and using (39) one last time, we finally get

$$\begin{aligned} \theta_{ur}^{\alpha_k}(\lambda) \theta_{vr}^{\alpha_k}(\lambda) &\leq 4M^3 r^{2d} \exp\left[-\frac{c}{8}(\alpha_k r)^\gamma\right] \pi_r^{\alpha_k}(\lambda) \\ &\stackrel{(42)}{\leq} \frac{\alpha_k^{3d}}{c_1 2^k} \pi_r^{\alpha_k}(\lambda), \end{aligned}$$

which concludes the proof using (40). In the second line, we used that R is chosen so large that for every $k \geq 0$ and $r \geq 3^k R$,

$$\frac{c_1 2^k}{\alpha_k^{3d}} 4M^3 r^{2d} \exp[-\frac{c}{8}(\alpha_k r)^\gamma] \leq \frac{c_1}{\alpha^{3d}} 4M^3 r^{6d} \exp[-\frac{c}{8}(\alpha r^{1-\bar{\gamma}})^\gamma] \leq 1. \quad (42)$$

□

Remark 23. *By adapting the reasoning above, one can prove similar statements for other distributions μ having sub-exponential tails, for instance $\mu[r, \infty]$ decaying like $\exp[-c(\log r)^\gamma]$ or $\exp[-cr/(\log r)^\gamma]$.*

Acknowledgments This research was supported by an IDEX grant from Paris-Saclay, the ERC grant CriBLaM, and the NCCR SwissMAP.

References

- [AB87] M. Aizenman and D. J. Barsky. Sharpness of the phase transition in percolation models. *Comm. Math. Phys.*, 108(3):489–526, 1987.
- [ABF87] M. Aizenman, D. J. Barsky, and R. Fernández. The phase transition in a general class of Ising-type models is sharp. *J. Statist. Phys.*, 47(3-4):343–374, 1987.
- [ATT16] D. Ahlberg, V. Tassion, and A. Teixeira. Sharpness of the phase transition for continuum percolation in \mathbb{R}^2 . *arXiv preprint arXiv:1605.05926*, 2016.
- [ATT17] D. Ahlberg, V. Tassion, and A. Teixeira. Existence of an unbounded vacant set for subcritical continuum percolation. *arXiv preprint arXiv:1706.03053*, 2017.
- [BD12] V. Beffara and H. Duminil-Copin. The self-dual point of the two-dimensional random-cluster model is critical for $q \geq 1$. *Probab. Theory Related Fields*, 153(3-4):511–542, 2012.
- [BH57] S. R. Broadbent and J. M. Hammersley. Percolation processes. I. Crystals and mazes. *Proc. Cambridge Philos. Soc.*, 53:629–641, 1957.
- [BR06] B. Bollobás and O. Riordan. The critical probability for random Voronoi percolation in the plane is $1/2$. *Probab. Theory Related Fields*, 136(3):417–468, 2006.
- [DGRSY18] H. Duminil-Copin, S. Goswami, A. Raoufi, F. Severo and A. Yadin, Existence of phase transition for percolation using the Gaussian Free Field, *arXiv:1806.07733*.

- [DGRS19] H. Duminil-Copin, S. Goswami, P.-F. Rodriguez and F. Severo, Equality of critical parameters for GFF level-set percolation, *in preparation*.
- [DRT17a] H. Duminil-Copin, A. Raoufi, and V. Tassion. Exponential decay of connection probabilities for subcritical Voronoi percolation in \mathbb{R}^d . To appear in *Probab. Theory Related Fields*. arXiv:1705.07978, 2017.
- [DRT17b] H. Duminil-Copin, A. Raoufi, and V. Tassion. Sharpness of the phase transition for random-cluster and Potts models via decision trees. arXiv:1705.03104, 2017.
- [DT16] H. Duminil-Copin and V. Tassion. A new proof of the sharpness of the phase transition for Bernoulli percolation and the Ising model. *Comm. Math. Phys.*, 343(2):725–745, 2016.
- [Gil61] E. N. Gilbert. Random plane networks. *J. Soc. Indust. Appl. Math.*, 9:533–543, 1961.
- [Gou08] J-B. Gou  r  . Subcritical regimes in the Poisson Boolean model of continuum percolation. *Ann. Probab.*, 36(4):1209–1220, 2008.
- [GT18] J-B. Gou  r   and M. Th  ret. Equivalence of some subcritical properties in continuum percolation, *arXiv:1803.00793*, 2018.
- [Men86] M. V. Menshikov. Coincidence of critical points in percolation problems. *Dokl. Akad. Nauk SSSR*, 288(6):1308–1311, 1986.
- [Hal85] P. Hall. On continuum percolation. *Ann. Probab.*, 13(4):1250–1266, 1985.
- [MR08] R. Meester and R. Roy. *Continuum percolation*. New York: Cambridge University Press, 2008.
- [MRS94] R. Meester, R. Roy, and A. Sarkar. Nonuniversality and continuity of the critical covered volume fraction in continuum percolation. *Journal of statistical physics*, 75(1-2):123–134, 1994.
- [Pen17] M. Penrose. Non-triviality of the vacancy phase transition for the Boolean model, arXiv:1706.02197, 2017.
- [OSSS05] R. O’Donnell, M. Saks, O. Schramm, and R. Servedio. Every decision tree has an influential variable. *FOCS*, 2005.
- [Zie16] S. Ziesche. Sharpness of the phase transition and lower bounds for the critical intensity in continuum percolation on \mathbb{R}^d , *to appear in AIHP*, arXiv:1607.06211, 2016.
- [ZS85] S. A Zuev and A. Sidorenko. Continuous models of percolation theory. I. *Theoretical and Mathematical Physics*, 62(1):51–58, 1985.