

The Riemann-Roch theorem for general elliptic operators

Mikhaël GROMOV and Mikhail A. SHUBIN

Abstract — The classical Riemann-Roch theorem is generalized to the solution of general elliptic equations with isolated singularities on an arbitrary compact manifold.

Le théorème de Riemann-Roch pour les opérateurs elliptiques généraux

Résumé — Le théorème de Riemann-Roch classique est généralisé pour les solutions d'équations elliptiques générales avec singularités isolées sur une variété compacte arbitraire.

Version française abrégée — Soient X une variété C^∞ fermée, $\dim X = n \geq 2$, E et F des fibrés vectoriels sur X , $q = \dim_{\mathbb{C}} E_x = \dim_{\mathbb{C}} F_x$ la dimension de leurs fibres, $\Gamma(U, E)$ l'espace des sections C^∞ de E sur un ouvert $U \subset X$. Soit $A : \Gamma(X, E) \rightarrow \Gamma(X, F)$ un opérateur elliptique différentiel d'ordre d , $\text{ind } A$ son indice qui peut être calculé par le théorème de l'indice d'Atiyah-Singer. Soient $\Omega(X)$ le fibré des densités complexes sur X , $E^* = \text{Hom}(E, \Omega(X))$, $\langle \cdot, \cdot \rangle$ la dualité bilinéaire naturelle $\Gamma(X, E) \times \Gamma(X, E^*) \rightarrow \mathbb{C}$ et A^t l'opérateur transposé défini par les dualités $\langle \cdot, \cdot \rangle$ dans E et F , ainsi $A^t : \Gamma(X, F^*) \rightarrow \Gamma(X, E^*)$ et

$$\langle Au, v \rangle = \langle u, A^t v \rangle, \quad u \in \Gamma(X, E), \quad v \in \Gamma(X, F^*).$$

Introduisons un diviseur $\mu = x_1^{p_1} x_2^{p_2} \dots x_m^{p_m}$, $x_i \in X$, $x_i \neq x_j$ si $i \neq j$, $p_i \in \mathbb{Z} \setminus \{0\}$, $\mu^{-1} = x_1^{-p_1} x_2^{-p_2} \dots x_m^{-p_m}$ est le diviseur dual. Le degré de μ est défini par

$$d(\mu) = \sum_{1 \leq i \leq m} \text{sign } p_i \left[\binom{|p_i| + n - 1}{n} - \binom{|p_i| + n - 1 - d}{n} \right]$$

où $\binom{N}{n} = N! / n!(N-n)!$ si $N \geq n$ et 0 sinon.

Soient $\text{supp } \mu = \{x_1, \dots, x_m\}$, $\lambda = \prod_{p_i \geq 0} x_i^{p_i}$, $\nu = \prod_{p_i \leq 0} x_i^{p_i}$. Soient

$L(\mu, A) = \{u \mid u \in \Gamma(X - \text{supp } \nu, E), Au = 0 \text{ dans } X - \text{supp } \nu;$

$$u(x) = o(|x - x_i|^{d-n-|p_i|}) \text{ lorsque } x \rightarrow x_i, x_i \in \text{supp } \nu;$$

$$u(x) = O(|x - x_i|^{p_i}) \text{ lorsque } x \rightarrow x_i, x \in \text{supp } \lambda \},$$

$$r(\mu, A) = \dim_{\mathbb{C}} L(\mu, A).$$

THÉORÈME 1. — $r(\mu, A) = \text{ind } A - qd(\mu) + r(\mu^{-1}, A^t)$.

Le théorème de Riemann-Roch classique est un cas particulier si X est une courbe algébrique complexe non-singulière, $A = \bar{\partial} : C^\infty(X) \rightarrow \Lambda^{0,1}(X)$. Le cas de l'opérateur de Laplace scalaire sur une variété riemannienne a été considéré par N. S. Nadirashvili [1].

Donnons quelques applications simples :

COROLLAIRE 3. — $r(\mu, A) \geq \text{ind } A - qd(\mu)$.

PROPOSITION 4. — Supposons que A^t a la propriété de prolongement unique, c'est-à-dire si $A^t U = 0$ et u a un zéro d'ordre infini en x_0 , alors $u = 0$ dans un voisinage de x_0 . Fixons $x_1, \dots, x_m \in X$ et $N_0 > 0$.

Note présentée par Mikhaël GROMOV.

Alors il existe $N > 0$ tel que, si $\sum_{p_i > 0} p_i \leq N_0$ et $\sum_{p_i < 0} |p_i| \geq N$, alors

$$r(\mu, A) = \text{ind } A - qd(\mu).$$

PROPOSITION 5. — Soit A une matrice $(q \times q)$ opérateur elliptique d'ordre d définie dans un voisinage de 0 dans \mathbb{R}^n . Soit $\mathcal{E}_0^{(k)}$ l'espace des germes de fonctions $f: U \rightarrow \mathbb{C}^q$ (U est un voisinage de 0 dans \mathbb{R}^n) telles que $\partial^d f(0) = 0$ pour $|d| \leq k$.

Alors l'application $A = \mathcal{E}_0^{(k+d)} \rightarrow \mathcal{E}_0^{(k)}$ est surjective.

1. INTRODUCTION. — The classical Riemann-Roch theorem for non-singular complex algebraic curves has been generalized in different ways to n -dimensional manifolds. The most known generalizations are the Riemann-Roch-Hirzebruch theorem and the Riemann-Roch-Grothendieck theorem in the algebraic geometry. In this Note we suggest a generalization motivated by the theory of solutions of elliptic equations with point singularities. The case of the scalar Laplacian was considered in a beautiful paper by N. S. Nadirashvili [1], which we took as a starting point. But our proof is simpler than that in [1] due to a use of duality arguments. We also give a number of applications including those which are similar to those of the classical Riemann-Roch theorem, and a local solvability result with an additional condition on the order of zero of the solution at the given point. Further applications will be given in a subsequent detailed paper.

2. NOTATIONS AND THE MAIN RESULT. — Let X be a compact C^∞ -manifold, $\dim X = n \geq 2$, E and F complex vector bundles over X , $q = \dim_{\mathbb{C}} E_x = \dim_{\mathbb{C}} F_x$ the dimension of their fibres, $\Gamma(U, E)$ the space of the C^∞ -sections of E over an open set $U \subset X$, $A: \Gamma(X, E) \rightarrow \Gamma(X, F)$ an elliptic differential operator of order d , $\text{ind } A$ its index which can be calculated by the Atiyah-Singer index formula, $\Omega(X)$ the bundle of complex densities on X , $E^* = \text{Hom}(E, \Omega(X))$, $\langle \cdot, \cdot \rangle$ the natural bilinear pairing $\Gamma(X, E) \times \Gamma(X, E^*) \rightarrow \mathbb{C}$ and A' the transposed operator to A defined by the pairings $\langle \cdot, \cdot \rangle$ in E and F , i.e. $A': \Gamma(X, F^*) \rightarrow \Gamma(X, E^*)$ and

$$\langle Au, v \rangle = \langle u, A'v \rangle, \quad u \in \Gamma(X, E), \quad v \in \Gamma(X, F^*).$$

Let us introduce a divisor $\mu = x_1^{p_1} x_2^{p_2} \dots x_m^{p_m}$, $x_i \in X$, $x_i \neq x_j$ if $i \neq j$, $p_i \in \mathbb{Z} \setminus \{0\}$, and $\mu^{-1} = x_1^{-p_1} x_2^{-p_2} \dots x_m^{-p_m}$ is the dual divisor. The degree of μ is defined as

$$d(\mu) = \sum_{i \leq m} \text{sign } p_i \left[\binom{|p_i| + n - 1}{n} - \binom{|p_i| + n - 1 - d}{n} \right]$$

where $\binom{N}{n} = N!/n!(N-n)!$ if $N \geq n$ and 0 otherwise. Let $\text{supp } \mu = \{x_1, \dots, x_m\}$, $\lambda = x_1^{p_1^+} \dots x_m^{p_m^+}$, $\nu = x_1^{p_1^-} \dots x_m^{p_m^-}$, where $p_i^+ = \max(p_i, 0)$, $p_i^- = \min(p_i, 0)$, so λ and ν are positive and negative part of μ respectively, $\text{supp } \mu = \text{supp } \lambda \cup \text{supp } \nu$, $\text{supp } \lambda \cap \text{supp } \nu = \emptyset$.

Now let us introduce the space of solutions of the equation $Au = 0$ with the prescribed point singularities ("poles") on $\text{supp } \nu$ and zeros of prescribed orders on $\text{supp } \lambda$:

$$L(\mu, A) = \{u \mid u \in \Gamma(X - \text{supp } \nu, E), Au = 0 \text{ on } X - \text{supp } \nu;$$

$$u(x) = o(|x - x_i|^{d-n-|p_i|}) \text{ as } x \rightarrow x_i, x_i \in \text{supp } \nu;$$

$$u(x) = O(|x - x_i|^{p_i}) \text{ as } x \rightarrow x_i, x \in \text{supp } \lambda \}.$$

The latter condition can be also written as $j_{x_i}^{p_i-1} u = 0, x \in \text{supp } \lambda$, where $j_{x_i}^{p_i-1} u$ is the jet of order p_i-1 of the section u at x_i . Now let $r(\mu, A) = \dim_{\mathbb{C}} L(\mu, A)$.

THEOREM 1:

$$(1) \quad r(\mu, A) = \text{ind } A - qd(\mu) + r(\mu^{-1}, A')$$

Examples. - (a) Let X be a compact connected complex manifold, $\dim_{\mathbb{C}} X = 1$, i.e. X is a complex non-singular algebraic curve. Take $A = \bar{\partial}: C^\infty(X) \rightarrow \Lambda^{0,1}(X)$ and identify $\Omega(X) = \Lambda^2(X)$ using the canonical orientation. Then $A' = \bar{\partial}: \Lambda^{1,0}(X) \rightarrow \Lambda^2(X)$ and $\text{ind } A = 1 - g$ where g is the genus of X . Theorem 1 then becomes the classical Riemann-Roch theorem if we note that here $n = 2, d = 1$, hence $d(\mu) = \sum_{1 \leq i \leq m} p_i$.

(b) Let X be a compact Riemannian manifold and $A = \Delta$ is the scalar Laplace-Beltrami operator. Then $A' = A$ and denoting $r(\mu) = r(\mu, \Delta)$ we obtain

$$r(\mu) = -d(\mu) + r(\mu^{-1}).$$

This was proved in [1] where also non-compact manifolds with boundary were allowed (with suitable boundary conditions and conditions at infinity).

3. PROOF OF THEOREM 1. - Denote $\mathcal{D}'(X, F)$ the space of distribution sections of F (dual space to $\Gamma(X, F^*)$) and $S(\mu, F)$ the space of $s \in \mathcal{D}'(X, F)$, such that $\text{supp } s \subset \text{supp } v$ and locally near the points $x_i \in \text{supp } v$ (i.e. the points entering in μ with negative exponents) s can be written as

$$s = \sum_{(i|p_i < 0)} \sum_{|\alpha| \leq |p_i| - 1} C_{i\alpha} \partial_x^\alpha \delta(x - x_i)$$

where δ is the Dirac measure, $C_{i\alpha} \in F_{x_i}$. So actually $S(\mu, F) = S(v, F)$.

Introduce the space $\Gamma(X, \mu, A)$ of sections $u \in \Gamma(X - \text{supp } v, E)$ having prescribed values and zeros [as in definition of $L(\mu, A)$] but not necessarily solutions of $Au = 0$:

$$\Gamma(X, \mu, A) = \{ u \mid u \in \Gamma(X - \text{supp } v, E), j_{x_i}^{p_i-1} u = 0 \text{ if } x_i \in \text{supp } \lambda;$$

for every $x_i \in \text{supp } v$ there exists a neighbourhood U of x_i

and a representation $u = u_s + u_r$ where $u_s \in \Gamma(U \setminus x_i, E)$,

$$Au = 0 \text{ in } U \setminus x_i, u(x) = O(|x - x_i|^{d-n-|p_i|}) \text{ as } x \rightarrow x_i,$$

and u_r can be extended to a section $\tilde{u}_r \in \Gamma(U, E)$ }.

For every $u \in \Gamma(X, \mu, A)$ we can find a "regularization" $\tilde{u} \in \mathcal{D}'(X, E)$ such that $\tilde{u} = u$ on $X - \text{supp } v$ and $A\tilde{u} = f + s$ with $f \in \Gamma(X, F)$ and $s \in S(\mu, F)$. Using the standard elliptic regularity result and the well-known structure of fundamental solutions of elliptic operators [2] it is easy to check that the space $\tilde{\Gamma}(X, \mu, A)$ of all such regularizations \tilde{u} can be described as a set of $\tilde{u} \in \mathcal{D}'(X, E)$ such that $\tilde{u} \in C^\infty$ is a neighbourhood of $\text{supp } \lambda$, $j_{x_i}^{p_i-1} u = 0$ for every $x_i \in \text{supp } \lambda$ and $A\tilde{u} = f + s$ with $f \in \Gamma(X, F)$ and $s \in S(\mu, F)$.

Now introduce the "reduced" divisor

$$\tilde{u} = x_1^{\tilde{p}_1} x_2^{\tilde{p}_2} \dots x_m^{\tilde{p}_m},$$

where $\tilde{p}_i = \text{sign } p_i \cdot (|p_i| - d)_+$ (the factors $x_i^{\tilde{p}_i}$ with $\tilde{p}_i = 0$ have to be omitted) and denote

$$\Gamma_{\tilde{\mu}}(X, F) = \{ f \mid f \in \Gamma(X, F), j_{x_i}^{\tilde{p}_i} f = 0 \text{ if } \tilde{p}_i > 0 \}.$$



