

ANALYTICITY OF THE SUSCEPTIBILITY FUNCTION
FOR UNIMODAL MARKOVIAN MAPS OF THE INTERVAL.

by Yunping Jiang* and David Ruelle**.

Abstract. We study the expression (susceptibility)

$$\Psi(\lambda) = \sum_{n=0}^{\infty} \lambda^n \int_I \rho(dx) X(x) \frac{d}{dx} A(f^n x)$$

where f is a unimodal Markovian map of the interval I , and $\rho = \rho_f$ is the corresponding absolutely continuous invariant measure. We show that $\Psi(\lambda)$ is analytic near $\lambda = 1$, where $\Psi(1)$ is formally the derivative of $\int_I \rho(dx) A(x)$ with respect to f in the direction of the vector field X .

* Math. Dept., Queens College of CUNY, Flushing, NY 11367 and Math. Dept., Graduate Center of CUNY and Math. Inst., AMSS, CAS. email: yunqc@forbin.qc.edu

** Math. Dept., Rutgers University, and IHES. 91440 Bures sur Yvette, France. email: ruelle@ihes.fr

In a previous note [Ru] the susceptibility function was analyzed for some examples of maps of the interval. The purpose of the present note is to give a concise treatment of the general unimodal Markovian case (assuming f real analytic). We hope that it will similarly be possible to analyze maps satisfying the Collet-Eckmann condition. Eventually, as explained in [Ru], application of a theorem of Whitney [Wh] should prove differentiability of the map $f \mapsto \rho_f$ restricted to a suitable set.

Setup

Let I be a compact interval of \mathbf{R} and $f : I \rightarrow I$ be real analytic. We assume that there is c in the interior of I such that $f'(0) = 0$, $f'(x) > 0$ for $x < c$, $f'(x) < 0$ for $x > c$, and $f''(c) < 0$. Replacing I by a possibly smaller interval, we assume that $I = [a, b]$ where $a = f^2(c)$, $b = f(c)$. We assume that the postcritical orbit $P = \{f^n c : n \geq 1\}$ is finite: $P = \{p_1, \dots, p_m\}$; in particular, f is Markovian. We shall assume that f is *analytically expanding* in the sense of Assumption A below; in particular the periodic orbits of f are assumed to be repelling, and therefore c cannot be periodic. We also assume that f is topologically mixing [this can always be achieved by replacing I by a smaller interval and f by some iterate f^N].

Theorem.

Under the above conditions, and Assumption A stated later, there is a unique f -invariant probability measure ρ absolutely continuous with respect to Lebesgue on I . If X is real analytic on I , and $A \in \mathbf{C}^1(I)$, then

$$\Psi(\lambda) = \sum_{n=0}^{\infty} \lambda^n \int_I \rho(dx) X(x) \frac{d}{dx} A(f^n x)$$

extends to a meromorphic function in \mathbf{C} , without pole on $\{\lambda : |\lambda| = 1\}$.

Change of variable

The finite set $\{c\} \cup P$ decomposes I into m subintervals I_j , with $2m$ endpoints (we “double” the endpoints of consecutive subintervals, distinguishing between a $-$ endpoint at the right of an interval, and a $+$ endpoint at the left). Note that $\eta = \{I_j : j = 1, \dots, m\}$ is a Markov partition for the map f . Consider the critical values of f^n . Then for large $n > 0$, the set of critical values will be stabilized and is always P . We define *polar* endpoints as follows:

(1) $p \in P$ is a polar $-$ endpoint of an interval in η if p is local maximum value of f^n for n large.

(2) $v \in P$ is a polar $+$ endpoint of an interval in η if p is local minimum value of f^n for n large.

Every $p \in P$ is a polar $-$ or $+$ endpoint and may be both, c is a nonpolar endpoint on both sides.

We define now an increasing continuous map $\varpi : I \rightarrow \mathbf{R}$ so that $J = \varpi I$ is a compact interval. We write $\varpi I_j = J_j$ for $1 \leq j \leq m$. Denote by ω the inverse of ϖ . We assume that $\omega|_{J_j}$ extends to a holomorphic function in a complex neighborhood of J_j for $1 \leq j \leq m$ and that for $q \in \{c\} \cup P$, ω has the property

$$\omega(\varpi q \pm \xi) = \omega(\varpi q) \pm \frac{\xi^2}{2} + O(\xi^4)$$

if q is a \pm polar endpoint, and

$$\omega(\varpi q \pm \xi) = \omega(\varpi q) \pm \xi + O(\xi^2)$$

if q is a nonpolar endpoint. [We should really consider disjoint copies of the I_j and J_j , and disjoint neighborhoods of these in \mathbf{C} or in a Riemann surface two-sheeted near polar endpoints. This would lead to notational complications that we prefer to omit].

Applications of this singular change of coordinate have been used in [Ji1], [BJR], and [Ru]; the reference [Ji2] contains some more relevant study regarding the method of singular change of coordinates in one-dimensional dynamical systems. The reader is encouraged to compare this method with orbifold metrics in [Th, Chapter 13]. Another relevant application of this method in complex dynamical systems can be found in [DH].

From now on we shall say that ϖq is a \pm polar (nonpolar) endpoint if q is \pm polar (nonpolar).

The dynamical system viewed after the change of variable.

For any two intervals $I_j, I_k \in \eta$ with $fI_j \supset I_k$, we define

$$\psi_{jk} = \varpi \circ (f|_{I_j})^{-1} \circ (\omega|_{J_k})$$

Note that the ψ_{jk} are restrictions of inverse branches of $g = \varpi \circ f \circ \omega : J \rightarrow J$ to intervals in η . The function $\psi_{jk} : J_k \rightarrow J_j$ extends holomorphically to a complex neighborhood of J_k . Indeed, note that $(f|_{I_j})^{-1}$ is holomorphic except if I_j is one of the two intervals around c , in which case the singularity is corrected by $\omega|_{I_n}$, where I_n is the rightmost interval in η . In other cases $\omega|_{I_k}$ cancels the singularity of $\varpi|_{I_j}$ by our definition of ω . [Note that $\psi'_{jk}(x) \geq 0$ or ≤ 0 on J_k and may vanish only at an interval endpoint].

Assumption A.

Each J_k , for $k = 1, \dots, m$, has a bounded open connected neighborhood U_k in \mathbf{C} such that $\psi_{jk} : J_k \rightarrow J_j$ extends to a continuous function $\psi_{jk} : \bar{U}_k \rightarrow \mathbf{C}$ holomorphic in U_k , and with $\psi_{jk}\bar{U}_k \subset U_j$.

One checks that the sets U_k can be assumed to be in ϵ -neighborhoods of the J_k . Also, Assumption A implies that periodic points for g are strictly repelling. The smoothness of ω , ϖ in the interior of subintervals shows that the same property holds for f , apart from interval endpoints where we however also assume the property to hold:

The periodic orbits of f are strictly repelling.

Markovian graph.

Consider the Markov partition $\eta = \{I_j\}$. Let us write $j \succ k$ (j covers k) if $fI_j \supset I_k$ (we allow $j \succ j$). This defines a directed graph with vertex set $\{1, \dots, m\}$ and oriented edges (j, k) for $j \succ k$. Since we have assumed our dynamical system f to be topological mixing, our graph is also mixing in the sense that there is $N \geq 1$ such that for all $j, k \in \{1, \dots, m\}$ we have $j \succ \dots \succ k$ (N edges) corresponding to $f^N I_j \supset I_k$.

Transfer operators.

For a function $\Phi = (\Phi_j)$ defined on $\sqcup J_j$, write

$$(\mathcal{L}\Phi)_k(z) = \sum_{j:j \succ k} \text{sgn}(j) \psi'_{jk}(z) \Phi(\psi_{jk}z)$$

$$(\mathcal{L}_0\Phi)_k(z) = \sum_{j:j \succ k} \text{sgn}(j) \Phi(\psi_{jk}z)$$

where $\text{sgn}(j)$ is $+1$ if ψ_{jk} is increasing on J_k , and -1 if ψ_{jk} is decreasing on J_k . If H is the Hilbert space of functions on $\sqcup_{j \in L} \bar{U}_j$ which are square integrable with respect to Lebesgue, and have holomorphic restrictions to the U_j , then \mathcal{L} and \mathcal{L}_0 acting on H are holomorphy improving, hence compact and of trace class.

Properties of \mathcal{L} .

For $x \in J_k$ we have

$$(\mathcal{L}\Phi)_k(x) = \sum_{j:j \succ k} |\psi'_{jk}(x)| \Phi_j(\psi_{jk}x)$$

hence $\Phi \geq 0$ implies $\mathcal{L}\Phi \geq 0$ (\mathcal{L} preserves positivity) and

$$\int_J dx (\mathcal{L}\Phi)(x) = \sum_k \int_{J_k} dx (\mathcal{L}\Phi)_k(x) = \sum_j \int_{J_j} dx \Phi_j(x) = \int_J dx \Phi(x)$$

(\mathcal{L} preserves mass). Using mixing one finds that \mathcal{L} has a simple eigenvalue $\mu_0 = 1$ corresponding to an eigenfunction $\sigma_0 > 0$. The other eigenvalues μ_ℓ satisfy $|\mu_\ell| < 1$, and their (generalized) eigenfunctions σ_ℓ satisfy $\int_J dx \sigma_\ell(x) = 0$. If we normalize σ_0 by $\int_J dx \sigma_0(x) = 1$, then $\sigma_0(dx) = \sigma_0(x)dx$ is the unique g -invariant probability measure absolutely continuous with respect to Lebesgue on J . In particular, $\sigma_0(x)dx$ is ergodic.

Let now $H_1 \subset H$ consist of those $\Phi = (\Phi_k)$ such that the derivative Φ' vanishes at the (polar) endpoints $\varpi a, \varpi b$ of J , and such that at the common endpoint ϖq ($q \in \{c\} \cup P \setminus \{a, b\}$) of two subintervals we have equality on both sides of a quantity which is either

- the value of Φ for a nonpolar endpoint, or
- the value of $\pm \Phi'$ for a polar \pm endpoint.

We note that $\mathcal{L}H_1 \subset H_1$ [this requires a case by case discussion]. Furthermore $\sigma_0 \in H_1$ [take $\phi \in H$ such that $\phi \geq 0$, $\int_J dy \phi(y) = 1$, and ϕ, ϕ' vanish at subinterval endpoints; then $\phi \in H_1$ and $\sigma_0 = \lim_{n \rightarrow \infty} \mathcal{L}^n \phi \in H_1$].

Evaluating $\Psi(\lambda)$.

The image $\rho(dx) = \rho(x)dx$ of $\sigma_0(y)dy$ by ω is the unique f -invariant probability measure absolutely continuous with respect to Lebesgue on I . We have

$$\rho(x) = \sigma_0(\varpi x) \varpi'(x)$$

Consider now the expression

$$\Psi(\lambda) = \sum_{n=0}^{\infty} \lambda^n \int_I \rho(dx) X(x) \frac{d}{dx} A(f^n x)$$

where we assume that X extends to a holomorphic function in a neighborhood of each I_k and takes the same value at both sides of common endpoints of intervals in η (continuity). Also assume that $A \in \mathcal{C}^1(I)$. For sufficiently small $|\lambda|$, the series defining $\Psi(\lambda)$ converges. Writing $B = A \circ \omega$ (B has piecewise continuous derivative) and $x = \omega y$ we have

$$X(x) \frac{d}{dx} A(f^n x) = X(\omega y) \frac{1}{\omega'(y)} \frac{d}{dy} B(g^n y)$$

hence

$$\Psi(\lambda) = \sum_{n=0}^{\infty} \lambda^n \int_J dy \sigma_0(y) \frac{X(\omega y)}{\omega'(y)} \frac{d}{dy} B(g^n y)$$

Defining $Y(y) = \sigma_0(y) X(\omega y) / \omega'(y)$, we see that Y extends to a holomorphic function in a neighborhood of each J_k , which we may take to be U_k , except for a simple pole at each polar endpoint of J_k . Since $\sigma_0 \in H_1$, the properties assumed for ω imply that also $(X \circ \omega) \times \sigma_0 \in H_1$. Note that near a nonpolar subinterval endpoint ϖq

$$\omega'(\varpi q \pm \xi) = 1 + O(\xi)$$

and near a \pm polar endpoint

$$\omega'(\varpi q \pm \xi) = \xi + O(\xi^3)$$

Therefore

$$Y(\varpi q \pm \xi) = A^\pm \frac{1}{\xi} + B^\pm + O(\xi)$$

where $B^+ = B^-$ for the two sides of ϖq , and $B^+ = 0$ at the left endpoint ϖa of J , $B^- = 0$ at the right endpoint ϖb of J . We may write

$$\int_J dy \sigma_0(y) \frac{X(\omega y)}{\omega'(y)} \frac{d}{dy} B(g^n y) = \int_J dy Y(y) g'(y) \cdots g'(g^{n-1} y) B'(g^n y)$$

$$= \int_J ds (\mathcal{L}_0^n Y)(s) B'(s)$$

where \mathcal{L}_0 has been defined above, and we have thus

$$\Psi(\lambda) = \sum_{n=0}^{\infty} \lambda^n \int_J ds (\mathcal{L}_0^n Y)(s) B'(s)$$

Properties of \mathcal{L}_0 .

We let now $H_0 \subset H$ be the space of functions $\Phi = (\Phi_k)$ vanishing at the endpoints $\varpi a, \varpi b$ of J , and such that the values of Φ on both sides of common endpoints of intervals J_j coincide (continuity). Therefore $\mathcal{L}_0 H_0 \subset H_0$.

There is a periodic orbit $\gamma_1, \dots, \gamma_p$ (with $g\gamma_j = \gamma_{j+1 \pmod{p}}$) of polar endpoints where γ_α is the \pm endpoint of some subinterval $J_{k(\alpha)}$. Choose P_α to be 0 on subintervals different from $J_{k(\alpha)}$, and to be holomorphic on a complex neighborhood of $J_{k(\alpha)}$ except at γ_α . Also assume that

$$P_\alpha(\gamma_\alpha \pm \xi) = \frac{1}{\xi} + O(\xi)$$

and that P_α vanishes at the endpoint of $J_{k(\alpha)}$ different from γ_α . Then

$$\mathcal{L}_0 P_\alpha - |f'(\gamma(\alpha))|^{1/2} P_{\alpha+1 \pmod{p}} \in H_0$$

Therefore $\mathcal{L}_0^p P_1 - \Lambda P_1 = u \in H_0$ where $\Lambda = \prod_{\alpha=1}^p |f'(\gamma(\alpha))|^{1/2} > 1$. Since the spectrum of \mathcal{L} acting on H is contained in the closed unit disk, and since the derivative u' is in H , we may define $v = (\mathcal{L}^p - \Lambda)^{-1} u' \in H$. Since $\int_J dy u'(y) = 0$ we also have $\int_J dy v(y) = 0$ and we can take $w \in H_0$ such that $w' = v$. We have thus

$$((\mathcal{L}_0^p - \Lambda)w)' = (\mathcal{L}^p - \Lambda)w' = (\mathcal{L}^p - \Lambda)v = u'$$

so that $(\mathcal{L}_0^p - \Lambda)w = u$ [there is no additive constant of integration since $(\mathcal{L}_0^p - \Lambda)w$ and u are in H_0]. Finally

$$(\mathcal{L}_0^p - \Lambda)(P_1 - w) = 0$$

There is thus a \mathcal{L}_0 -invariant p -dimensional vector space spanned by vectors $P_\alpha - w_\alpha$ with $w_\alpha \in H_0$, such that the spectrum of \mathcal{L}_0 restricted to this space consists of eigenvalues ω_ℓ with

$$\omega_\ell = \Lambda^{1/p} e^{2\pi \ell i / p} = \left| \prod_{\alpha=1}^p f'(\gamma_\alpha) \right|^{1/2p} e^{2\pi \ell i / p}$$

for $\ell = 0, \dots, p-1$.

For the postcritical but nonperiodic polar points $\tilde{\gamma}_1, \dots, \tilde{\gamma}_q$ define \tilde{P}_β like P_α above, with γ_α replaced by $\tilde{\gamma}_\beta$. For each β there is $\alpha = \alpha(\beta)$ with

$$\mathcal{L}_0^q(\tilde{P}_\beta - \tilde{\Lambda}_\beta P_\alpha) \in H_0$$

with some $\tilde{\Lambda}_\beta \neq 0$, hence

$$\mathcal{L}_0^q(\tilde{P}_\beta - \tilde{\Lambda}_\beta(P_\alpha - w_\alpha)) = \tilde{Y}_\beta \in H_0$$

Poles of $\Psi(\lambda)$.

We may now write

$$Y = Y_0 + Y_1 + Y_2$$

where

$$\begin{aligned} Y_0 &\in H_0 \\ Y_1 &= \sum_{\alpha=1}^p c_\alpha(P_\alpha - w_\alpha) \\ Y_2 &= \sum_{\beta=1}^q \tilde{c}_\beta(\tilde{P}_\beta - \tilde{\Lambda}_\beta(P_{\alpha(\beta)} - w_{\alpha(\beta)})) \end{aligned}$$

and there is a corresponding decomposition $\Psi(\lambda) = \Psi_0(\lambda) + \Psi_1(\lambda) + \Psi_2(\lambda)$. Here $\Psi_1(\lambda)$ is a sum of terms $C_\ell/(\lambda - \omega_\ell)$ where $\omega_\ell = \Lambda^{1/p} \times p$ -th root of unity; $\Psi_2(\lambda) =$ polynomial of degree $q - 1$ in λ plus $\lambda^q \sum_{\beta=1}^q \tilde{c}_\beta \tilde{\Psi}_\beta(\lambda)$ where $\tilde{\Psi}_\beta$ is obtained if we replace Y by \tilde{Y}_β in the definition of Ψ . The poles of $\Psi(\lambda)$ are thus those of $\Psi_1(\lambda)$ at the ω_ℓ and those of $\Psi_0(\lambda)$ and $\tilde{\Psi}_\beta(\lambda)$. The discussion is the same for Ψ_0 and the $\tilde{\Psi}_\beta$, we shall thus only consider Ψ_0 . Since $Y_0 \in H_0$ and $\mathcal{L}_0 H_0 \subset H_0$ we have

$$\begin{aligned} \Psi_0(\lambda) &= \sum_{n=0}^{\infty} \lambda^n \int_J ds (\mathcal{L}_0^n Y_0)(s) B'(s) = - \sum_{n=0}^{\infty} \lambda^n \int_J ds (\mathcal{L}_0^n Y_0)'(s) B(s) \\ &= - \sum_{n=0}^{\infty} \lambda^n \int_J ds (\mathcal{L}^n Y_0')(s) B(s) \end{aligned}$$

It follows that $\Psi_0(\lambda)$ extends meromorphically to \mathbf{C} with poles at the μ_k^{-1} . We want to show that the residue of the pole at $\mu_0^{-1} = 1$ vanishes. Since $\int_J dy \sigma_k(y) = 0$ for $k \geq 1$, the coefficient of σ_0 in the expansion of Y_0' is proportional to

$$\int_J dy Y_0'(y) = Y(\varpi b) - Y(\varpi a) = 0$$

because $Y_0 \in H_0$. Therefore $\Psi_0(\lambda)$ is holomorphic for $|\lambda| = 1$, and the same holds for the $\tilde{\Psi}_\beta(\lambda)$, concluding the proof of the theorem. In fact we know that the poles of $\Psi(\lambda)$ are located at μ_k^{-1} for $k \geq 1$, and at ω_ℓ^{-1} for $\ell = 0, \dots, p - 1$, so that $|\mu_k^{-1}| > 1, |\omega_\ell^{-1}| < 1$.

Acknowledgments.

One of us (Y.J.) is partially supported by grants from the NSF and the PSC-CUNY Program, and by the Hundred Talents Program of the CAS.

References.

- [BJR] V. Baladi, Y. Jiang, H.H. Rugh. “Dynamical determinants via dynamical conjugacies for postcritically finite polynomials.” *J. Statist. Phys.* **108**, 973-993(2002).
- [DH] D. Douady and J. H. Hubbard, “A proof of Thurston’s topological characterization of rational functions.” *Acta Math.*, Vol. **171**, 263-297(1993).
- [Ji1] Y. Jiang, “On Ulam-von Neumann transformations.” *Commun. Math. Phys.*, **172**, 449-459(1995).
- [Ji2] Y. Jiang, “Renormalization and Geometry in One-Dimensional and Complex Dynamics.” *Advanced Series in Nonlinear Dynamics*, Vol. 10 (1996), World Scientific Publishing Co. Pte. Ltd., River Edge, NJ.
- [Ru] D. Ruelle. “Differentiating the absolutely continuous invariant measure of an interval map f with respect to f .” *Commun. Math. Phys.* to appear.
- [Th] W. Thurston, “The Geometry and Topology of Three-Manifolds”, Electronic version 1.1-March 2002, <http://www.msri.org/publications/books/gt3m/>
- [Wh] H. Whitney. “Analytic expansions of differentiable functions defined in closed sets.” *Trans. Amer. Math. Soc.* **36**,63-89(1934).