

PERTURBATION THEORY FOR LYAPUNOV EXPONENTS OF A TORAL MAP:
EXTENSION OF A RESULT OF SHUB AND WILKINSON.

by David Ruelle*.

Abstract. Starting from a hyperbolic toral automorphism times a rotation of the circle, we obtain, for a small volume preserving perturbation, an exact and rigorous second order perturbation expansion of the Lyapunov exponents.

Keywords: Lyapunov exponent, toral automorphism, hyperbolicity.

* Mathematics Dept., Rutgers University, and IHES. 91440 Bures sur Yvette, France.
<ruelle@ihes.fr>

We consider volume preserving perturbations F of a diffeomorphism $F_0 = (\Phi, J)$ of $\mathbf{T}^{m+1} = \mathbf{T}^m \times \mathbf{T}$, where Φ is a hyperbolic automorphism of \mathbf{T}^m , and J is a translation of \mathbf{T} . Writing $F = F_0 + aF'$, we shall show that the Lyapunov exponents for (F, volume) can be expanded to second order in a (Theorem 1). In particular, the central Lyapunov exponent λ^c of (F, volume) , to second order in a , is generally $\neq 0$ (Corollary 11). For a special family of perturbations one obtains particularly simple formulae, first noted by Shub and Wilkinson [17]. We recover their result in Theorem 12. We deviate from [17] mostly in that we don't have differentiability of λ^c , only a second order expansion around $a = 0$. The ideas used here are largely those in Shub and Wilkinson [17], and can be appreciated in the background provided by Hirsh, Pugh and Shub [9], Burns and Wilkinson [5], Ruelle and Wilkinson [16], Nițică and Török [12], Pugh, Shub and Wilkinson [14]. Among older regularity results let us mention Katok, Knieper and Weiss [11], Flaminio [9], Ruelle [15]. For recent work concerning Lyapunov exponents, see Bonatti, Gómez-Mont and Viana [3], Avila and Bochi [2]. Closely related to the subject of the present paper are the references [4] and [6].

After completing the writing of this paper, the author received a preprint by D. Dolgopyat [7], which develops similar ideas in a more general setting, but without the specific formulas we obtain here.

1. Theorem.

Let Φ be a hyperbolic automorphism of \mathbf{T}^m , and $J : y \mapsto y + \alpha \pmod{1}$ a translation of \mathbf{T} . Define $F_0 = (\Phi, J)$, and let $F = F_0 + aF'$ (+higher order in a) be a C^2 perturbation of F_0 , volume preserving to first order in a . (We take $F' : \mathbf{T}^{m+1} \mapsto \mathbf{R}^{m+1}$ and $F_0\xi + aF'(\xi)$ has to be understood $\pmod{1}$ in each component). Let $\lambda_1 < \lambda_2 < \dots$ be the Lyapunov exponents of (F_0, volume) and m_1, m_2, \dots their multiplicities (the exponent = 0 occurs with multiplicity 1). Also let $\lambda_a^{(1)} \leq \lambda_a^{(2)} \leq \dots$ be the Lyapunov exponents of (F, volume) repeated according to multiplicity. Then we have the second order expansion

$$\sum_{\ell=m_1+\dots+m_{r-1}+1}^{m_1+\dots+m_r} \lambda_a^{(\ell)} = m_r \lambda_r + a^2 L_r + o(a^2)$$

If $m_r = 1$, and writing $\lambda_r = \lambda_0^{(\ell)}$, we have

$$\lambda_a^{(\ell)} = \lambda_0^{(\ell)} + a^2 L^{(\ell)} + o(a^2)$$

(this applies in particular to $\lambda^c = \lambda_a^{(\ell)}$ for $\lambda_0^{(\ell)} = 0$).

An explicit expression for L_r can be obtained (see Proposition 9). We do not assume ergodicity of (F, volume) , and therefore we use *integrated* Lyapunov exponents (averaged over the volume), see however Remark 15(a).

Because the perturbation $+aF'$ (+higher order in a) to F_0 gives only a quadratic contribution in the above formulas, the higher order terms do not contribute to order a^2 . Since the higher order terms do not change our results, these terms will be omitted in what follows.

2. Normal hyperbolicity.

As in [17], we invoke the theory of normal hyperbolicity of [10]. We start from the fact that F_0 is normally hyperbolic to the smooth fibration of \mathbf{T}^{m+1} by circles $\{x\} \times \mathbf{T}$. Taking some $k \geq 2$ we apply [10] Theorems (7.1), (7.2). Thus we obtain a C^1 neighborhood U of F_0 in the C^k diffeomorphisms of \mathbf{T}^{m+1} such that, for $F \in U$, there is an equivariant fibration $\pi : \mathbf{T}^{m+1} \rightarrow \mathbf{T}^m$ with

$$\pi F = \Phi \pi$$

The fibers $\pi^{-1}\{x\}$ are C^k circles forming a continuous fibration of \mathbf{T}^{m+1} (this fibration is in general not smooth). Furthermore there is a TF -invariant continuous splitting of $T\mathbf{T}^{m+1}$ into three subbundles:

$$T\mathbf{T}^{m+1} = E^s + E^u + E^c$$

such that E^c is 1-dimensional tangent to the circles $\pi^{-1}\{x\}$, E^s is m^s -dimensional contracting and E^u is m^u -dimensional expanding for TF .

If $\lambda_r < 0$ (and F is in a suitable C^1 -small neighborhood U of F_0), we can introduce a continuous vector subbundle E^r of $T\mathbf{T}^{m+1}$ which consists of vectors contracting under TF^n faster than $(\lambda_r + \epsilon)^n$ where $\epsilon > 0$ and $\lambda_r + \epsilon < \lambda_{r+1}$. In fact E^r is a hyperbolic (attracting) fixed point for the action induced by TF^{-1} on the bundle of $m_1 + \dots + m_r$ dimensional linear subspaces of $T\mathbf{T}^{m+1}$ (over F^{-1} acting on T^{m+1}).

If $\lambda_r > 0$, replacement of F by F^{-1} similarly yields a continuous subbundle \bar{E}^r of $m_r + \dots$ dimensional subspaces.

3. Proposition.

Assume that F is of class C^k , $k \geq 2$, and that F is C^k close to F_0 . The bundles E^r , \bar{E}^r , when restricted to a circle $\pi^{-1}\{x\}$ are of class C^{k-1} , continuously in x .

If \mathcal{G} denotes the (Grassmannian) manifold of $m_1 + \dots + m_r$ dimensional linear subspaces of \mathbf{R}^{m+1} , we may identify the bundle of $m_1 + \dots + m_r$ dimensional linear subspaces of $T\mathbf{T}^{m+1}$ with $\mathbf{T}^{m+1} \times \mathcal{G}$. We denote by $\mathcal{E} \in \mathcal{G}$ the spectral subspace of the matrix defining Φ corresponding to the smallest $m_1 + \dots + m_r$ eigenvalues (in absolute value, and repeated according to multiplicity).

If \mathcal{F}_0 is the action defined by TF_0 on $T\mathbf{T}^{m+1} \times \mathcal{G}$, the circles $\{x\} \times \mathbf{T} \times \{\mathcal{E}\}$ form an \mathcal{F}_0 invariant fibration of $\mathbf{T}^{m+1} \times \{\mathcal{E}\}$, to which \mathcal{F}_0 is normally hyperbolic. If F is C^k close to F_0 , the corresponding C^{k-1} action \mathcal{F} is normally hyperbolic to a perturbed fibration where $\{x\} \times \mathbf{T} \times \{\mathcal{E}\}$ is replaced by $E^r|_{\pi^{-1}\{x\}}$. According to [10] Theorem 7.4, Corollary (8.3) and the following Remark 2, we find that the C^{k-1} circle $E^r|_{\pi^{-1}\{x\}} \subset \mathbf{T}^{m+1} \times \mathcal{G}$ depends continuously on $x \in \mathbf{T}^{m+1}$. Similarly for \bar{E} . \square

Note that in [17], the C^r section theorem is used in a similar situation, giving estimates uniform in x . However, continuity in x (not just uniformity) will be essential for us in what follows.

4. Corollary.

The splitting $T\mathbf{T}^{m+1} = E^s + E^u + E^c$ when restricted to a circle $\pi^{-1}\{x\}$ is of class C^{k-1} , continuously in x .

It is clear that $E^c|_{\pi^{-1}\{x\}}$ is of class C^{k-1} because it is the tangent bundle to the C^k circle $\pi^{-1}\{x\}$. As to E^s, E^u , they are special cases of E^r, \bar{E}^r . \square

Notation.

Remember that $F = F_0 + aF'$, and fix F' . We shall use the notation π_a, E_a^r, \dots to indicate the a -dependence of π, E^r, \dots

5. Proposition.

For small $\epsilon > 0$ there is a continuous function $x \mapsto \gamma_x$ from \mathbf{T}^m to $C^k(\mathbf{T} \times (-\epsilon, \epsilon) \rightarrow \mathbf{T}^m)$ such that $\gamma_x(y, 0) = 0$ and $\pi_a^{-1}\{x\} = \{(x + \gamma_x(y, a), y) : y \in \mathbf{T}\}$.

To see this define $\tilde{F} : \mathbf{T}^{m+1} \times (-\epsilon, \epsilon) \rightarrow \mathbf{T}^{m+1} \times (-\epsilon, \epsilon)$ by $\tilde{F}(\xi, a) = ((F_0 + aF')(\xi), a)$ and observe that \tilde{F} is normally hyperbolic to the 2-dimensional manifolds

$$\cup_{a \in (-\epsilon, \epsilon)} (\pi_a^{-1}\{x\}, a)$$

and these are thus C^k 2-dimensional submanifolds of $\mathbf{T}^{m+1} \times (-\epsilon, \epsilon)$. \square

We may in the same manner replace $\pi_a^{-1}\{x\}$ by $\cup_{a \in (-\epsilon, \epsilon)} (\pi_a^{-1}\{x\}, a)$ in Proposition 3 and Corollary 4. Writing E_a for $E_a^r, \bar{E}_a^r, E_a^s, E_a^u, E_a^c$, we obtain that $(\cdot, a) \mapsto E_a(\cdot)$, when restricted from $\mathbf{T}^{m+1} \times (-\epsilon, \epsilon)$ to $\cup_{a \in (-\epsilon, \epsilon)} (\pi_a^{-1}\{x\}, a)$ is of class C^{k-1} . We rephrase this as follows:

6. Proposition.

The map

$$x \mapsto \{(y, a) \mapsto E_a(x + \gamma_x(y, a), y)\}$$

where E_a stands for $E_a^r, \bar{E}_a^r, E_a^s, E_a^u, E_a^c$, is continuous $\mathbf{T}^m \rightarrow C^{k-1}(\mathbf{T} \times (-\epsilon, \epsilon) \rightarrow \text{Grassmannian of } \mathbf{R}^{m+1})$ where we have used the identification $T\mathbf{T}^{m+1} = \mathbf{T}^{m+1} \times \mathbf{R}^{m+1}$. \square

Notation.

From now on we write E_a for $E_a^r, \bar{E}_a^r, E_a^s, E_a^u, E_a^c$. When $a = 0$, E_0 is a trivial subbundle of $T\mathbf{T}^{m+1} = \mathbf{T}^{m+1} \times \mathbf{R}^{m+1}$, and we shall write $E_0 = \mathbf{T}^{m+1} \times \mathcal{E}$, denoting thus by \mathcal{E} a spectral subspace of the matrix on \mathbf{R}^{m+1} defining $(\Phi, 1)$. We denote by \mathcal{E}^\perp the complementary spectral subspace.

Taking $k = 2$ we have then:

7. Corollary.

There are linear maps $G(x, y), R(x, y, a) : \mathcal{E} \rightarrow \mathcal{E}^\perp$ such that $G(x, y)$ depends continuously on $(x, y) \in \mathbf{T}^m \times \mathbf{T}$, $R(x, y, a)$ on $(x, y, a) \in \mathbf{T}^m \times \mathbf{T} \times (-\epsilon, \epsilon)$,

$$E_a(x + \gamma_x(y, a), y) = \{X + aG(x, y)X + R(x, y, a)X : X \in \mathcal{E}\}$$

and $\|R(x, y, a)\|$ is $o(a)$ uniformly in x, y . \square

Notice now that, if $\tilde{x} = \pi_a(x, y)$, then $x = \tilde{x} + \gamma_{\tilde{x}}(y, a)$, where $\gamma_{\tilde{x}}(y, a) = O(a)$. Now

$$E_a(x, y) = E_a(\tilde{x} + \gamma_{\tilde{x}}(y, a), y) = \{X + aG(\tilde{x}, y)X + R(\tilde{x}, y, a)X : X \in \mathcal{E}\}$$

differs from

$$E_a(x + \gamma_x(y, a), y) = \{X + aG(x, y)X + R(x, y, a)X : X \in \mathcal{E}\}$$

by the replacement $\tilde{x} \rightarrow x$ in the right-hand side, and since $\text{dist}(\tilde{x}, x) = O(a)$, we find that $\text{dist}(E_a(x, y), E_a(x + \gamma_x(y, a), y)) = o(a)$. Therefore, changing the definition of R , we can again write:

8. Corollary.

There are linear maps $G(x, y), R(x, y, a) : \mathcal{E} \rightarrow \mathcal{E}^\perp$, depending continuously on their arguments, such that

$$E_a(x, y) = \{X + aG(x, y)X + R(x, y, a)X : X \in \mathcal{E}\}$$

and $\|R(x, y, a)\|$ is $o(a)$ uniformly in x, y . \square

We may write $T_\xi F = T_\xi(F_0 + aF') = D_0 + aD'(\xi)$ where D_0 does not depend on ξ and preserves the decomposition $T_\xi M = \mathcal{E} + \mathcal{E}^\perp$. If we apply TF to an element $X + aGX + RX$ of E_a (as in Corollary 8) we obtain $X_1 +$ element of $\mathcal{E}^\perp \in E_a$, with $X_1 \in \mathcal{E}$:

$$X_1 = D_0X + aD'X + a^2D'GX + aD'RX \quad \text{projected on } \mathcal{E} \quad (1)$$

Under $(TF)^\wedge$, the volume element θ in $E_a(\xi)$ is multiplied by a factor $M(\xi, a)$, and the projection in \mathcal{E} of $(TF)^\wedge\theta$ is equal to the projection in \mathcal{E} of θ multiplied by a factor $N(\xi, a)$ such that

$$M(\xi, a) = N(\xi, a) + \ell_a(\xi) - \ell_a(F\xi)$$

for suitable ℓ_a . We may compute N from (1):

$$N(\xi, a) = N_{(0)} + aN_{(1)}(\xi) + a^2N_{(2)}(\xi) + o(a^2)$$

To proceed we take now $E_a = E_a^r$, and assume $\lambda_r < 0$. We have then, writing $d\xi$ for the volume element in \mathbb{T}^{m+1} ,

$$\begin{aligned} L_a &= \sum_{\ell=1}^{m_1+\dots+m_r} \lambda_a^{(\ell)} = \int d\xi \log M(\xi, a) = \int d\xi \log N(\xi, a) \\ &= L_{(0)} + aL_{(1)}(\xi) + a^2L_{(2)}(\xi) + o(a^2) \end{aligned} \quad (2)$$

More precisely, we shall prove

9. Proposition.

If $\lambda_r < 0$, we have

$$\sum_{\ell=1}^{m_1+\dots+m_r} \lambda_a^{(\ell)} = \sum_{k=1}^r m_k \lambda_k + a^2 L + o(a^2)$$

where

$$L = \frac{1}{2} \sum_{n=-\infty}^{\infty} \int d\xi \operatorname{Tr}_{\mathcal{E}}(D_0^{-1} D'(\xi)) \operatorname{Tr}_{\mathcal{E}}(D_0^{-1} D'(F_0^n \xi)) \geq 0$$

and $\operatorname{Tr}_{\mathcal{E}}$ is defined as follows. Let \mathcal{E} be the spectral subspace of the matrix D_0 (defining $(\Phi, 1)$ in \mathbf{R}^{m+1}) corresponding to the smallest $m_1 + \dots + m_r$ eigenvalues (in absolute value, and repeated according to multiplicity). Also let \mathcal{E}^{\perp} be the complementary spectral subspace. We define P to be the projection on \mathcal{E} parallel to \mathcal{E}^{\perp} , and write $\operatorname{Tr}_{\mathbf{R}^{m+1}} P \dots P$.

The convergence of the series defining L is exponential as will result from the proof. We postpone showing that $L \geq 0$ until Remark 15(b).

The proposition is obtained by comparing formula (2) with the formula (5) below, which we shall obtain by a second order perturbation calculation.

To first order in a we have

$$F^n = (F_0 + aF')^n = F_0^n + a \sum_{j=1}^n F_0^{n-j} \circ F' \circ F_0^{j-1}$$

hence

$$T_{\xi} F^n = D_0^n + a \sum_{j=1}^n D_0^{n-j} D'(F^{j-1} \xi) D_0^{j-1}$$

If we apply $T F^n$ to $X + aGX + RX \in E_a$ we obtain $X_n +$ element of $\mathcal{E}^{\perp} \in E_a$, with $X_n \in \mathcal{E}$. To zero-th order in a , $X_n = D_0^n X$, so we may write to first order $X_n = D_0^n X + aY_n(\xi)$. Therefore, to first order in a ,

$$D_0^n X + aY_n(\xi) + aG(F^n \xi) D_0^n X = D_0^n X + a \sum_{j=1}^n D_0^{n-j} D'(F^{j-1} \xi) D_0^{j-1} X + aD_0^n G(\xi) X$$

and, taking the components along \mathcal{E}^{\perp} ,

$$G(F^n \xi) D_0^n X = \sum_{j=1}^n D_0^{n-j} D'_{\perp}(F^{j-1} \xi) D_0^{j-1} X + D_0^n G(\xi) X$$

where $D'_{\perp}(\cdot)$ is $D'(\cdot)$ followed by taking the component along \mathcal{E}^{\perp} , or

$$\sum_{j=1}^n D_0^{-j} D'_{\perp}(F^{j-1} \xi) D_0^{j-1} X + G(\xi) X = D_0^{-n} G(F^n \xi) D_0^n X$$

When $n \rightarrow \infty$, the right-hand side tends to zero (exponentially fast, remember that $X \in \mathcal{E}$, $GX \in \mathcal{E}^\perp$). Therefore (to order 0 in a)

$$G(\xi)X = - \sum_{j=1}^{\infty} D_0^{-j} D'_\perp (F^{j-1} \xi) D_0^{j-1} X$$

which we shall use in the form

$$G(\xi)X = - \sum_{n=0}^{\infty} D_0^{-n-1} D'_\perp (F_0^n \xi) D_0^n X \quad (3)$$

where we have written F_0^n instead of F^n since G is evaluated to order 0 in a . (The right-hand side is an exponentially convergent series).

Returning to (1) we see that, to second order in a ,

$$\begin{aligned} X_1 &= D_0 X + a D'(\xi) X + a^2 D'(\xi) G(\xi) X \quad \text{projected on } \mathcal{E} \\ &= D_0 (1 + a D_0^{-1} D'(\xi) + a^2 D_0^{-1} D'(\xi) G(\xi)) X \quad \text{projected on } \mathcal{E} \end{aligned}$$

Let now $(u^{(i)})$ and $(u^{(i)\perp})$ be conjugate bases of \mathcal{E} . Also let $\delta^{(i)}$ for $i = 1, \dots, m_1 + \dots + m_r$ be the eigenvalues of D_0 restricted to \mathcal{E} . Then, to second order in a ,

$$N(\xi, a) \wedge_1^{m_1 + \dots + m_r} u^{(\ell)}$$

is, up to a factor of absolute value 1,

$$\begin{aligned} & \left(\prod_{\ell=1}^{m_1 + \dots + m_r} \delta^{(\ell)} \right) [1 + a \sum_{i=1}^{m_1 + \dots + m_r} (u^{(i)\perp}, D_0^{-1} D'(\xi) u^{(i)}) \\ & + a^2 \sum_{i < j} ((u^{(i)\perp}, D_0^{-1} D'(\xi) u^{(i)}) (u^{(j)\perp}, D_0^{-1} D'(\xi) u^{(j)}) \\ & - (u^{(i)\perp}, D_0^{-1} D'(\xi) u^{(j)}) (u^{(j)\perp}, D_0^{-1} D'(\xi) u^{(i)})) + a^2 \sum_i (u^{(i)\perp}, D_0^{-1} D'(\xi) G(\xi) u^{(i)})] \wedge_\ell u^{(\ell)} \end{aligned}$$

so that

$$\begin{aligned} N(\xi, a) &= \left(\prod_{\ell=1}^{m_1 + \dots + m_r} |\delta^{(\ell)}| \right) [1 + \{ a \sum_i (u^{(i)\perp}, D_0^{-1} D'(\xi) u^{(i)}) \\ & + a^2 \sum_{i < j} ((u^{(i)\perp}, D_0^{-1} D'(\xi) u^{(i)}) (u^{(j)\perp}, D_0^{-1} D'(\xi) u^{(j)}) \\ & - (u^{(i)\perp}, D_0^{-1} D'(\xi) u^{(j)}) (u^{(j)\perp}, D_0^{-1} D'(\xi) u^{(i)})) + a^2 \sum_i (u^{(i)\perp}, D_0^{-1} D'(\xi) G(\xi) u^{(i)}) \}] \end{aligned}$$

Since $\log |\delta^{(\ell)}| = \lambda_0^{(\ell)}$ we obtain, to second order in a ,

$$L_a = \int d\xi \log N(\xi, a) = m_1 \lambda_1 + \dots + m_r \lambda_r + \int d\xi [\{\dots\} - \frac{a^2}{2} (\sum_i (u^{(i)\perp}, D_0^{-1} D'(\xi) u^{(i)}))^2]$$

where $\{\dots\}$ has the same meaning as above. Write

$$\Psi_i(\sum_\ell \xi_\ell u^{(\ell)}) = (u^{(i)\perp}, D_0^{-1} F'(\sum_\ell \xi_\ell u^{(\ell)}))$$

The first term of $\int d\xi \{\dots\}$ is

$$a \sum_i \int d\xi (u^{(i)\perp}, D_0^{-1} T F'(\xi) u^{(i)}) = a \sum_i \int d\xi \frac{\partial}{\partial \xi_i} \Psi_i$$

which vanishes because $\int d\xi \frac{\partial}{\partial \xi_i} \dots = 0$. The next term in $\int d\xi \{\dots\}$ is

$$a^2 \sum_{i < j} \int d\xi \left(\left(\frac{\partial \Psi_i}{\partial \xi_i} \right) \left(\frac{\partial \Psi_j}{\partial \xi_j} \right) - \left(\frac{\partial \Psi_i}{\partial \xi_j} \right) \left(\frac{\partial \Psi_j}{\partial \xi_i} \right) \right) = a^2 \sum_{i < j} \int d\xi \left(\frac{\partial}{\partial \xi_i} (\Psi_i \frac{\partial \Psi_j}{\partial \xi_j}) - \frac{\partial}{\partial \xi_j} (\Psi_i \frac{\partial \Psi_j}{\partial \xi_i}) \right)$$

which vanishes as above. Thus we are left with

$$\begin{aligned} & L_a - (m_1 \lambda_1 + \dots + m_r \lambda_r) \\ &= a^2 \int d\xi \left[\sum_i (u^{(i)\perp}, D_0^{-1} D'(\xi) G(\xi) u^{(i)}) - \frac{1}{2} (\sum_i (u^{(i)\perp}, D_0^{-1} D'(\xi) u^{(i)}))^2 \right] \end{aligned} \quad (4)$$

and we may write, using (3),

$$\begin{aligned} \sum_i (u^{(i)\perp}, D_0^{-1} D'(\xi) G(\xi) u^{(i)}) &= - \sum_{n=0}^{\infty} \sum_i (u^{(i)\perp}, D_0^{-1} D'(\xi) D_0^{-n-1} D'_\perp (F_0^n) D_0^n u^{(i)}) \\ &= - \sum_{n=0}^{\infty} \sum_i \sum_j^* (u^{(i)\perp}, D_0^{-1} D'(\xi) u^{(j)}) (u^{(j)\perp}, D_0^{-n-1} D'(F_0^n \xi) D_0^n u^{(i)}) \end{aligned}$$

where we have introduced conjugate bases $(u^{(j)}), (u^{(j)\perp})$ of \mathcal{E}^\perp , indexed by $j = m_1 + \dots + m_r + 1, \dots, m + 1$, and \sum_i is over $i \leq m_1 + \dots + m_r$, \sum_j^* is over $j \geq m_1 + \dots + m_r + 1$. The above expression is also

$$= - \sum_{n=0}^{\infty} \sum_i \sum_j^* \frac{\partial}{\partial \xi_j} (u^{(i)\perp}, D_0^{-1} F'(\sum_\ell \xi_\ell u^{(\ell)})) \frac{\partial}{\partial \xi_i} (u^{(j)\perp}, D_0^{-n-1} F'(F_0^n \sum_\ell \xi_\ell u^{(\ell)}))$$

and integration by part gives thus

$$\int d\xi \sum_i (u^{(i)\perp}, D_0^{-1} D'(\xi) G(\xi) u^{(i)})$$

$$\begin{aligned}
&= - \sum_{n=0}^{\infty} \int d\xi \sum_i \frac{\partial}{\partial \xi_i} (u^{(i)\perp}, D_0^{-1} F'(\sum_{\ell} \xi_{\ell} u^{(\ell)})) \sum_j^* \frac{\partial}{\partial \xi_j} (u^{(j)\perp}, D_0^{-n-1} F'(F_0^n \sum_{\ell} \xi_{\ell} u^{(\ell)})) \\
&= - \sum_{n=0}^{\infty} \int d\xi \operatorname{Tr}_{\mathcal{E}}(D_0^{-1} D'(\xi)) \operatorname{Tr}_{\mathcal{E}^{\perp}}(D_0^{-n-1} D'(F_0^n \xi) D_0^n) \\
&= - \sum_{n=0}^{\infty} \int d\xi \operatorname{Tr}_{\mathcal{E}}(D_0^{-1} D'(\xi)) \operatorname{Tr}_{\mathcal{E}^{\perp}}(D_0^{-1} D'(F_0^n \xi))
\end{aligned}$$

(here $\operatorname{Tr}_{\mathcal{E}^{\perp}} = \operatorname{Tr}_{\mathbf{R}^{m+1}} - \operatorname{Tr}_{\mathcal{E}}$). The fact that $F = F_0 + aF'$ is volume preserving (to first order in a) is expressed by $\operatorname{Tr}_{\mathbf{R}^{m+1}}(D_0^{-1} D'(\xi)) = 0$ hence

$$\begin{aligned}
&\int d\xi \sum_i (u^{(i)\perp}, D_0^{-1} D'(\xi) G(\xi) u^{(i)}) \\
&= \sum_{n=0}^{\infty} \int d\xi \operatorname{Tr}_{\mathcal{E}}(D_0^{-1} D'(\xi)) \operatorname{Tr}_{\mathcal{E}}(D_0^{-1} D'(F_0^n \xi))
\end{aligned}$$

and introducing this in (4) yields

$$\begin{aligned}
&L_a - (m_1 \lambda_1 + \dots + m_r \lambda_r) \\
&= a^2 \left[\sum_{n=1}^{\infty} \int d\xi \operatorname{Tr}_{\mathcal{E}}(D_0^{-1} D'(\xi)) \operatorname{Tr}_{\mathcal{E}}(D_0^{-1} D'(F_0^n \xi)) + \frac{1}{2} \int d\xi (\operatorname{Tr}_{\mathcal{E}}(D_0^{-1} D'(\xi)))^2 \right] \\
&= \frac{a^2}{2} \sum_{n=-\infty}^{\infty} \int d\xi \operatorname{Tr}_{\mathcal{E}}(D_0^{-1} D'(\xi)) \operatorname{Tr}_{\mathcal{E}}(D_0^{-1} D'(F_0^n \xi)) \tag{5}
\end{aligned}$$

where the last step used the invariance of $d\xi$ under F_0^n . \square

10. Proof of Theorem 1.

We use Proposition 9, the corresponding result with F replaced by F^{-1} , and the fact that $\sum_{\ell=1}^m \lambda_a^{(\ell)} = 0$ (because F is volume preserving). This gives an estimate of all the sums of $\lambda_a^{(\ell)}$ that occur in Theorem 1. \square

11. Corollary.

In the situation of Theorem 1, the central Lyapunov exponent is

$$\begin{aligned}
\lambda^c &= \frac{a^2}{2} \sum_{-\infty}^{\infty} \int d\xi [\operatorname{Tr}^u(D_0^{-1} D'(\xi)) \operatorname{Tr}^u(D_0^{-1} D'(F_0^n \xi)) - \operatorname{Tr}^s(D_0^{-1} D'(\xi)) \operatorname{Tr}^s(D_0^{-1} D'(F_0^n \xi))] \\
&= \frac{a^2}{2} \sum_{-\infty}^{\infty} \int d\xi [\operatorname{Tr}^s(D_0^{-1} D'(\xi)) - \operatorname{Tr}^u(D_0^{-1} D'(\xi))] \operatorname{Tr}^c(D_0^{-1} D'(F_0^n \xi))
\end{aligned}$$

where Tr^s , Tr^u , Tr^c denote the traces over the spectral subspaces \mathcal{E}^s , \mathcal{E}^u , \mathcal{E}^c of D_0 corresponding to eigenvalues < 1 , > 1 , or $= 1$ in absolute value (\mathcal{E}^c is one dimensional).

Since F preserves the volume, the sum of all Lyapunov exponents vanishes. Therefore λ^c is minus the sum of the negative Lyapunov exponents, given by (5), minus the sum of the positive Lyapunov exponents. Note that replacing F by F^{-1} , \mathcal{E}^s by \mathcal{E}^u (and, to the order considered, $D'(\xi)$ by $-D'(\xi)$) replaces the sum of the negative Lyapunov exponents by minus the sum of the positive exponents. This gives the first formula for λ^c .

To obtain the second formula, express $\text{Tr}^u \text{Tr}^u - \text{Tr}^s \text{Tr}^s$ in terms of $\text{Tr}^u \pm \text{Tr}^s$, and remember that (because F preserves the volume) $\text{Tr}^s + \text{Tr}^u + \text{Tr}^c = 0$ when applied to $D_0^{-1} D'(\xi)$. \square

The above formula (5) takes a particularly simple form in a special case described in the next theorem.

12. Theorem.

Let Φ be a hyperbolic automorphism of \mathbf{T}^m , with stable and unstable dimensions m^s and $m^u = m - m^s$, and with entropy λ_0^u . Let $J : y \rightarrow y + \alpha \pmod{1}$ be a translation of \mathbf{T} , and $\phi : \mathbf{T}^m \rightarrow \mathbf{T}$ a group homomorphism $\neq 0$. Finally let $\psi : \mathbf{T} \rightarrow \mathbf{R}^m$ be a nullhomotopic C^2 function.

Define $h, g_a : \mathbf{T}^m \times \mathbf{T} \rightarrow \mathbf{T}^m \times \mathbf{T}$ by

$$h \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} \Phi x \\ Jy + \phi \Phi x - \phi x \end{pmatrix}, \quad g_a \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x + a\psi(y) \pmod{1} \\ y \end{pmatrix}$$

and let $f_a = g_a \circ h$.

Denote by λ_a^s (resp. λ_a^u) the sum of the smallest m^s (resp. the largest m^u) Lyapunov exponents for (f_a, volume) . Also let $\lambda_a^c = -\lambda_a^s - \lambda_a^u$ be the ‘‘central exponent’’. Then λ_a^s , λ_a^u , λ_a^c have expansions of order 2 in a :

$$\lambda_a^s = -\lambda_0^u + \frac{a^2}{2} \int_{\mathbf{T}} dy ((\nabla \phi) \psi'^s(y))^2 + o(a^2)$$

$$\lambda_a^u = \lambda_0^u - \frac{a^2}{2} \int_{\mathbf{T}} dy ((\nabla \phi) \psi'^u(y))^2 + o(a^2)$$

$$\lambda_a^c = \frac{a^2}{2} \int_{\mathbf{T}} dy [((\nabla \phi) \psi'^u(y))^2 - ((\nabla \phi) \psi'^s(y))^2] + o(a^2)$$

Here $\psi'^s(y)$ and $\psi'^u(y)$ are the components of the derivative $\psi'(y) \in \mathbf{R}^m$ in the stable and unstable subspaces \mathcal{E}^s and \mathcal{E}^u for Φ . Also, we have used $\nabla \phi : \mathbf{R}^m \rightarrow \mathbf{R}$ to denote the derivative of the map $\phi : \mathbf{T}^m \rightarrow \mathbf{T}$ with the obvious identifications.

This theorem is a simple (but nontrivial) extension of the result proved by Shub and Wilkinson [17]. In the situation that they consider $\Phi = \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix}$, $J = \text{identity}$, $\phi = (1, 0)$,

$\psi' = \psi'^u$. [Remark that, in the notation of [17], $u_0 = ((1, 1).v_0)/(m - 1) = ((1, 0).v_0)$ so that the formula given in Proposition II of [17] agrees with our result above].

Notation.

We shall henceforth omit the (mod 1). We shall keep ∇ to denote the derivative in \mathbf{T}^m . With obvious abuses of notation, the reader may find it convenient to think of Φ or $\nabla\Phi$ as an $m \times m$ matrix (with integer entries and determinant ± 1), and ϕ or $\nabla\phi$ as a row m -vector (with integer entries not all zero).

13. Reformulation of the problem.

Note that $f_a^{-1} = h^{-1} \circ g_a^{-1}$ where h^{-1}, g_a^{-1} are obtained from h, g_a by the replacements $\Phi, J, \phi, \psi \rightarrow \Phi^{-1}, J^{-1}, \phi, -\psi$. These replacements also interchange the stable and unstable subspaces for Φ and replace λ^s, λ^u by $-\lambda^u, -\lambda^s$. Therefore the formula for λ^u in the theorem follows from the formula for λ^s . And the formula for $\lambda^c = -\lambda^s - \lambda^u$ also follows. To complete the proof of the theorem we turn now to the formula for λ^s .

Define

$$\hat{\phi}\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \\ y + \phi x \end{pmatrix}$$

then

$$F_0\begin{pmatrix} x \\ y \end{pmatrix} = \hat{\phi}^{-1}h\hat{\phi}\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} \Phi x \\ Jy \end{pmatrix}$$

$$\hat{g}_a\begin{pmatrix} x \\ y \end{pmatrix} = \hat{\phi}^{-1}g_a\hat{\phi}\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x + a\psi(y + \phi x) \\ y - a(\nabla\phi)\psi(y + \phi x) \end{pmatrix}$$

so that

$$F\begin{pmatrix} x \\ y \end{pmatrix} = \hat{\phi}^{-1}f_a\hat{\phi}\begin{pmatrix} x \\ y \end{pmatrix} = \hat{g}_a F_0\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} \Phi x + a\psi(Jy + \phi\Phi x) \\ Jy - a(\nabla\phi)\psi(Jy + \phi\Phi x) \end{pmatrix}$$

Finally, $F = F_0 + aF'$ with

$$F_0\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} \Phi x \\ Jy \end{pmatrix}, \quad F'\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} \psi(Jy + \phi\Phi x) \\ -(\nabla\phi)\psi(Jy + \phi\Phi x) \end{pmatrix}$$

Since F is conjugate (linearly) to f_a , we may compute λ^s from F instead of f_a .

14. Proof of Theorem 12.

Write $\mathbf{R}^{m+1} = \mathcal{E}^s + \mathcal{E}^u + \mathbf{R}$. We shall apply Proposition 9 with $\mathcal{E} = \mathcal{E}^s$, $\mathcal{E}^\perp = \mathcal{E}^u + \mathbf{R}$. Using $\xi = (x, y)$ and $X \in \mathcal{E}^s$, $Y \in \mathcal{E}^u$, $Z \in \mathbf{R}$ we may write

$$D_0\begin{pmatrix} X + Y \\ Z \end{pmatrix} = \begin{pmatrix} (\nabla\Phi)(X + Y) \\ Z \end{pmatrix}$$

$$D'(\xi)\begin{pmatrix} X + Y \\ Z \end{pmatrix} = \begin{pmatrix} \psi'(Jy + \phi\Phi x)((\nabla\phi\Phi)(X + Y) + Z) \\ -(\nabla\phi)\psi'(Jy + \phi\Phi x)((\nabla\phi\Phi)(X + Y) + Z) \end{pmatrix}$$

where ψ' denotes the derivative of ψ . Therefore

$$\text{Tr}_{\mathcal{E}}(D'(\xi)D_0^{-1}) = (\nabla\phi)\psi'^s(Jy + \phi\Phi x)$$

and (5) contains the integrals

$$\begin{aligned} & \int d\xi \operatorname{Tr}_{\mathcal{E}}(D_0^{-1}D'(\xi))\operatorname{Tr}_{\mathcal{E}}(D_0^{-1}D'(F_0^n\xi)) \\ &= \int d\xi [(\nabla\phi)\psi'^s(Jy + \phi\Phi x)][(\nabla\phi)\psi'^s(J^{n+1}y + \phi\Phi^{n+1}x)] \end{aligned}$$

Performing a change of variables $\bar{x} = \Phi x$, $\bar{y} = Jy + \phi\Phi x$ we find that this is

$$= \int d\bar{x} d\bar{y} [(\nabla\phi)\psi'^s(\bar{y})][(\nabla\phi)\psi'^s(J^n\bar{y} + \phi\Phi^n\bar{x} - \phi\bar{x})]$$

We claim that this last integral vanishes unless $n = 0$. This is because, if $n \neq 0$,

$$\int d\bar{x} \psi'(J^n\bar{y} + \phi\Phi^n\bar{x} - \phi\bar{x}) = 0$$

Indeed, $\phi\Phi^n\bar{x} - \phi\bar{x}$ is a linear combination with integer coefficients of the components $\bar{x}_1, \dots, \bar{x}_m$ of \bar{x} , and the coefficients do not all vanish because $\phi\Phi^n = \phi$ is impossible (Φ is hyperbolic and $\phi \neq 0$). Integrating the derivative ψ' with respect to a variable \bar{x}_j really occurring in $\phi\Phi^n\bar{x} - \phi\bar{x}$ gives zero as announced.

Returning to (5) we have thus

$$\begin{aligned} \lambda_a^s + \lambda_0^u &= \frac{a^2}{2} \int d\xi (\operatorname{Tr}_{\mathcal{E}}(D_0^{-1}D'(\xi)))^2 \\ &= \frac{a^2}{2} \int d\bar{y} ((\nabla\phi)\psi'^s(\bar{y}))^2 \end{aligned}$$

which is the formula given for λ_a^s in Theorem 12. And according to Section 13 this completes our proof. \square

15. Final remarks.

(a) Shub and Wilkinson [17] showed that close to a diffeomorphism (hyperbolic automorphism Φ of \mathbf{T}^2) \times (identity on \mathbf{T}) there is a C^1 open set of ergodic volume preserving C^2 diffeomorphisms of \mathbf{T}^3 with central Lyapunov exponent $\lambda^c > 0$. They remark that their result extends to the situation where Φ is a hyperbolic automorphism of \mathbf{T}^m with one-dimensional expanding eigenspace. More generally, if Φ is any hyperbolic automorphism of \mathbf{T}^m , Theorem 12 gives close to $(\Phi, \text{rotation of } \mathbf{T})$ in $C^2(\mathbf{T}^{m+1})$ a diffeomorphism F with $\lambda^c > 0$. Since λ^c is given by an integral over the volume of a local ‘‘central’’ stretching exponent, we have $\lambda^c > 0$ in a C^1 neighborhood of F . But by a result of Dolgopyat and Wilkinson [8] (Corollary 0.5), stable ergodicity is here C^1 open and dense in the C^2 volume preserving diffeomorphisms (C^1 is improved to C^k in [12]): we have center bunching and stable dynamical coherence because we consider perturbations of $(\Phi, \text{rotation of } \mathbf{T})$ for which the center foliation is C^1 , see [10], [13]. In conclusion, close to (hyperbolic automorphism Φ of \mathbf{T}^m) \times (rotation on \mathbf{T}) there is a C^1 open set V of ergodic volume preserving C^2

diffeomorphisms of \mathbf{T}^{m+1} with central Lyapunov exponent $\lambda^c > 0$ (or also with $\lambda^c < 0$). In particular, if $F \in V$, the conditional measures of the volume on the circles $\pi^{-1}\{x\}$ are atomic, as discussed in [16].

(b) The coefficient L in Proposition 9 is ≥ 0 . Consider indeed the unitary operator U defined by $U\psi = \psi \circ F$ on $L^2(\mathbf{T}^{m+1}, \text{volume})$, and let $E(\cdot)$ be the corresponding spectral measure, so that

$$U = \int_{\mathbf{T}} e^{2\pi i\theta} E(d\theta)$$

If $\psi(\xi) = \text{Tr}_{\mathcal{E}}(D_0^{-1}D'(\xi))$ we have a measure $\nu \geq 0$ on \mathbf{T} defined by $\nu(d\theta) = (\psi, E(d\theta)\psi)$ and the Fourier coefficients

$$c_n = \int e^{2\pi ni\theta} \nu(d\theta) = \int d\xi \text{Tr}_{\mathcal{E}}(D_0^{-1}D'(\xi))(D_0^{-1}D'(F_0^n\xi))$$

of this measure tend to zero exponentially. Therefore $\nu(d\theta) = \rho(\theta)d\theta$ has a smooth density ρ and

$$L = \frac{1}{2} \sum_{n=-\infty}^{\infty} c_n = \frac{1}{2} \rho(0) \geq 0$$

(c) Suppose now that F is not necessarily a volume preserving perturbation of F_0 . We may still hope that F has an SRB measure ρ_a . If F_0 were hyperbolic, we would have an expansion

$$\rho_a = \rho_0 + a\delta + o(a)$$

(see [15]) with $\rho_0 = \text{Lebesgue measure}$ and δ a distribution. For smooth Ψ , $\delta(\Psi)$ is given (because ρ_0 is Lebesgue measure) by the simple formula (see [15])

$$\delta(\Psi) = - \sum_0^{\infty} \rho_0((\Psi \circ F_0^n) \cdot \text{div}(F' \circ F_0^{-1}))$$

Similarly (replacing F by F^{-1} , hence $F_0, D_0^{-1}D'(\xi)$ by $F_0^{-1}, -D'(F_0^{-1}\xi)D_0^{-1}$) we see that the anti-SRB state has an expansion

$$\bar{\rho}_a = \rho_0 + a\bar{\delta} + o(a)$$

with

$$\begin{aligned} \bar{\delta}(\Psi) &= \sum_{n=1}^{\infty} \int d\xi \Psi(F_0^{-n}\xi) \text{Tr}_{\mathbf{R}^{m+1}}(D'(F_0^{-1}\xi)D_0^{-1}) \\ &= \sum_{n=0}^{\infty} \int d\xi \Psi(F_0^{-n}\xi) \text{Tr}_{\mathbf{R}^{m+1}}(D_0^{-1}D'(\xi)) \end{aligned}$$

We can now estimate the Lyapunov exponents for (F, ρ_a) to second order in a even though we are not sure of the existence of the SRB measure ρ_a . We simply assume that we

can use the formula for $\delta(\Psi)$. Going through the proof of Proposition 9 we have to replace $\int d\xi \log N(\xi, a)$ by $\rho_a(\log N(\cdot, a))$ and (to second order in a) this adds to the right-hand side of (4) a term

$$-a^2 \sum_{n=1}^{\infty} \int d\xi \operatorname{Tr}_{\mathcal{E}}(D_0^{-1}D'(\xi)) \operatorname{Tr}_{\mathbf{R}^{m+1}}(D_0^{-1}D'(\xi))$$

Taking into account the integrations by part we obtain now instead of (5) the formula

$$\begin{aligned} L_a - (m_1 \lambda_1 + \dots + m_r \lambda_r) &= \frac{a^2}{2} \sum_{n=-\infty}^{\infty} \int d\xi \operatorname{Tr}_{\mathcal{E}}(D_0^{-1}D'(\xi)) \operatorname{Tr}_{\mathcal{E}}(D_0^{-1}D'(F_0^n \xi)) \\ &\quad - a^2 \sum_{n=-\infty}^{\infty} \int d\xi \operatorname{Tr}_{\mathcal{E}}(D_0^{-1}D'(\xi)) \operatorname{Tr}_{\mathbf{R}^{m+1}}(D_0^{-1}D'(F_0^n \xi)) \end{aligned} \quad (6)$$

Let $a^2 L^s$, $a^2 L^u$, $a^2 L^c$ be the a^2 contributions to the sum of the noncentral negative, noncentral positive, and the central Lyapunov exponents for the SRB measure. We obtain $a^2 L^s$ from (6) when $n_r = n^s$. A similar calculation gives $a^2 L^u$ (it is convenient here to work via the anti-SRB measure, then replace F by F^{-1}). Estimating the average expansion coefficient gives $a^2(L^s + L^u + L^c) = \rho_a(\log \det(D_0 + aD'(\cdot)))$, hence $L^s + L^u + L^c$, hence L^c . The results are

$$\begin{aligned} L^s &= \frac{1}{2} \sum_{n=-\infty}^{\infty} \int d\xi \operatorname{Tr}^s(D_0^{-1}D'(\xi)) \operatorname{Tr}^s(D_0^{-1}D'(F_0^n \xi)) \\ &\quad - \sum_{n=-\infty}^{\infty} \int d\xi \operatorname{Tr}^s(D_0^{-1}D'(\xi)) \operatorname{Tr}_{\mathbf{R}^{m+1}}(D_0^{-1}D'(F_0^n \xi)) \\ L^u &= -\frac{1}{2} \sum_{n=-\infty}^{\infty} \int d\xi \operatorname{Tr}^u(D_0^{-1}D'(\xi)) \operatorname{Tr}^u(D_0^{-1}D'(F_0^n \xi)) \\ L^c &= -\frac{1}{2} \sum_{n=-\infty}^{\infty} \int d\xi \operatorname{Tr}^c(D_0^{-1}D'(\xi)) \operatorname{Tr}^c(D_0^{-1}D'(F_0^n \xi)) \\ &\quad - \sum_{n=-\infty}^{\infty} \int d\xi \operatorname{Tr}^c(D_0^{-1}D'(\xi)) \operatorname{Tr}^u(D_0^{-1}D'(F_0^n \xi)) \\ L^s + L^u + L^c &= -\frac{1}{2} \sum_{n=-\infty}^{\infty} \int d\xi \operatorname{Tr}_{\mathbf{R}^{m+1}}(D_0^{-1}D'(\xi)) \operatorname{Tr}_{\mathbf{R}^{m+1}}(D_0^{-1}D'(F_0^n \xi)) \end{aligned}$$

which can be rewritten variously.

In view of recent work [4], [1], [6], it seems reasonable to conjecture that if the above L^c is $\neq 0$, then there exists an SRB measure for (small) finite a .

Acknowledgements.

I am indebted to Mike Shub, Marcelo Viana, Amie Wilkinson, and Lai-Sang Young for a number of useful conversations related to the present article. Also I wish to thank the referee for helpful remarks.

References.

- [1] J. Alves, C. Bonatti and M. Viana. “SRB measures for partially hyperbolic systems whose central direction is mostly expanding.” *Inventiones Math.* **140**,351-398(2000).
- [2] A. Avila and J. Bochi. “A formula with some applications to the theory of Lyapunov exponents.” Preprint.
- [3] C. Bonatti, X. Gómez-Mont and M. Viana. “Généricité d'exponents de Lyapunov non-nuls des produits déterministes de matrices.” Preprint.
- [4] C. Bonatti and M. Viana. “SRB measures for partially hyperbolic systems whose central direction is mostly contracting.” *Israel J. Math.* **115**,157-193(2000).
- [5] K. Burns and A. Wilkinson. “Stable ergodicity of skew products.” *Ann. Sci. Ecole Norm. Sup.* **32**,859-889(1999).
- [6] W. Cowieson and L.-S. Young. Work in progress.
- [7] D. Dolgopyat. “On differentiability of SRB states.” Preprint.
- [8] D. Dolgopyat and A. Wilkinson. “Stable accessibility is C^1 dense.” Preprint
- [9] L. Flaminio. “Local entropy rigidity for hyperbolic manifolds.” *Commun. Anal. Geom.* **3**,555-596(1995).
- [10] M. Hirsch, C. Pugh, and M. Shub. *Invariant manifolds*. Lect. Notes in Math. **583** Springer, Berlin, 1977.
- [11] A. Katok, G. Knieper and H. Weiss. “Formulas for the derivative and critical points of topological entropy for Anosov and geodesic flows.” *Commun. Math. Phys.* **138**,19-31(1991).
- [12] V. Nițică and A. Török. “An open dense set of stably ergodic diffeomorphisms in a neighborhood of a non-ergodic one.” Preprint.
- [13] C. Pugh and M. Shub. “Stable ergodicity and julienne quasi-conformality.” *J. Eur. Math. Soc.* **2**,1-52(2000).
- [14] C. Pugh, M. Shub and A. Wilkinson. “Differentiating E^u .” In preparation.
- [15] D. Ruelle. “Differentiation of SRB states.” *Commun. Math. Phys.* **187**,227-241(1997).
- [16] D. Ruelle and A. Wilkinson. “Absolutely singular dynamical foliations.” *Commun. Math. Phys.*, to appear.
- [17] M. Shub and A. Wilkinson. “Pathological foliations and removable exponents.” *Inventiones Math.* **139**,495-508(2000).