

Kinematics and Cohomology

Dirk Kreimer

March 18, 2010

Hopf algebra of graphs $H = \mathbb{Q}1 \oplus \bigoplus_{j=1}^{\infty} H^j$

► The coproduct

$$\Delta(\Gamma) = \Gamma \otimes 1 + 1 \otimes \Gamma + \overbrace{\sum_{\gamma = \cup_i \gamma_i, \omega_4(\gamma_i) \geq 0}}^{\Delta'(\Gamma)} \gamma \otimes \Gamma/\gamma \quad (1)$$

Hopf algebra of graphs $H = \mathbb{Q}1 \oplus \bigoplus_{j=1}^{\infty} H^j$

► The coproduct

$$\Delta(\Gamma) = \Gamma \otimes 1 + 1 \otimes \Gamma + \overbrace{\sum_{\gamma = \cup_i \gamma_i, \omega_4(\gamma_i) \geq 0} \gamma \otimes \Gamma/\gamma}^{\Delta'(\Gamma)} \quad (1)$$

► The antipode

$$S(\Gamma) = -\Gamma - \sum S(\gamma)\Gamma/\gamma = -m(S \otimes P)\Delta \quad (2)$$

Hopf algebra of graphs $H = \mathbb{Q}1 \oplus \bigoplus_{j=1}^{\infty} H^j$

► The coproduct

$$\Delta(\Gamma) = \Gamma \otimes 1 + 1 \otimes \Gamma + \overbrace{\sum_{\gamma = \cup_i \gamma_i, \omega_4(\gamma_i) \geq 0} \gamma \otimes \Gamma/\gamma}^{\Delta'(\Gamma)} \quad (1)$$

► The antipode

$$S(\Gamma) = -\Gamma - \sum S(\gamma)\Gamma/\gamma = -m(S \otimes P)\Delta \quad (2)$$

► The character group

$$G_V^H \ni \Phi \Leftrightarrow \Phi : H \rightarrow V, \Phi(h_1 \cup h_2) = \Phi(h_1)\Phi(h_2) \quad (3)$$

Hopf algebra of graphs $H = \mathbb{Q}1 \oplus \bigoplus_{j=1}^{\infty} H^j$

► The coproduct

$$\Delta(\Gamma) = \Gamma \otimes 1 + 1 \otimes \Gamma + \overbrace{\sum_{\gamma = \cup_i \gamma_i, \omega_4(\gamma_i) \geq 0} \Gamma \otimes \Gamma/\gamma}^{\Delta'(\Gamma)} \quad (1)$$

► The antipode

$$S(\Gamma) = -\Gamma - \sum S(\gamma)\Gamma/\gamma = -m(S \otimes P)\Delta \quad (2)$$

► The character group

$$G_V^H \ni \Phi \Leftrightarrow \Phi : H \rightarrow V, \Phi(h_1 \cup h_2) = \Phi(h_1)\Phi(h_2) \quad (3)$$

► The counterterm

$$\begin{aligned} S_R^\Phi(\Gamma) &= -R \left(\Phi(h) - \sum S_R^\Phi(\gamma)\Phi(\Gamma/\gamma) \right) \\ &= -R \Phi \left(m(S_R^\Phi \otimes \Phi P)\Delta(\Gamma) \right) \end{aligned} \quad (4)$$

Hopf algebra of graphs $H = \mathbb{Q}1 \oplus \bigoplus_{j=1}^{\infty} H^j$

► The coproduct

$$\Delta(\Gamma) = \Gamma \otimes 1 + 1 \otimes \Gamma + \overbrace{\sum_{\gamma = \cup_i \gamma_i, \omega_4(\gamma_i) \geq 0} \Gamma \otimes \Gamma/\gamma}^{\Delta'(\Gamma)} \quad (1)$$

► The antipode

$$S(\Gamma) = -\Gamma - \sum S(\gamma)\Gamma/\gamma = -m(S \otimes P)\Delta \quad (2)$$

► The character group

$$G_V^H \ni \Phi \Leftrightarrow \Phi : H \rightarrow V, \Phi(h_1 \cup h_2) = \Phi(h_1)\Phi(h_2) \quad (3)$$

► The counterterm

$$\begin{aligned} S_R^\Phi(\Gamma) &= -R \left(\Phi(h) - \sum S_R^\Phi(\gamma)\Phi(\Gamma/\gamma) \right) \\ &= -R \Phi \left(m(S_R^\Phi \otimes \Phi P)\Delta(\Gamma) \right) \end{aligned} \quad (4)$$

► The renormalized Feynman rules

$$\Phi_R = m(S_R^\Phi \otimes \Phi)\Delta \quad (5)$$

An Example

- ▶ The co-product

$$\Delta' \left(\begin{array}{c} \text{Diagram 1} \\ \text{Diagram 2} \\ \text{Diagram 3} \\ \text{Diagram 4} \\ \text{Diagram 5} \\ \text{Diagram 6} \\ \text{Diagram 7} \end{array} \right) = 3 \begin{array}{c} \text{Diagram 8} \\ \otimes \\ \text{Diagram 9} \end{array} + 2 \begin{array}{c} \text{Diagram 10} \\ \otimes \\ \text{Diagram 11} \end{array} + \begin{array}{c} \text{Diagram 12} \\ \otimes \\ \text{Diagram 13} \end{array}.$$

An Example

- ▶ The co-product

$$\Delta' \left(\text{diagram} \right) = 3 \text{diagram} \otimes \text{diagram} + 2 \text{diagram} \otimes \text{diagram} + \text{diagram} \otimes \text{diagram} .$$

- ▶ The counterterm

$$\begin{aligned} S_R^\Phi \left(\text{diagram} \right) &= -Rm \left[S_R^\Phi \otimes \Phi P \right] \times \\ &\quad \times \Delta \left(\text{diagram} \right) \\ &= -R \left\{ \Phi \left(\text{diagram} \right) + \right. \\ &\quad \left. + R \left[\Phi \left(3 \text{diagram} + 2 \text{diagram} + \text{diagram} \right) \right] \Phi \left(\text{diagram} \right) \right\} \end{aligned}$$

An Example

- ▶ The co-product

$$\Delta' \left(\text{diagram} \right) = 3 \text{diagram} \otimes \text{diagram} + 2 \text{diagram} \otimes \text{diagram} + \text{diagram} \otimes \text{diagram} .$$

- ▶ The counterterm

$$\begin{aligned} S_R^\Phi \left(\text{diagram} \right) &= -Rm \left[S_R^\Phi \otimes \Phi P \right] \times \\ &\quad \times \Delta \left(\text{diagram} \right) \\ &= -R \left\{ \Phi \left(\text{diagram} \right) + \right. \\ &\quad \left. + R \left[\Phi \left(3 \text{diagram} + 2 \text{diagram} + \text{diagram} \right) \right] \Phi \left(\text{diagram} \right) \right\} \end{aligned}$$

- ▶ The renormalized result

$$\begin{aligned} \Phi_R &= (\text{id} - R)m(S_R^\Phi \otimes \Phi P)\Delta \left(\text{diagram} \right) \\ &= (\text{id} - R) \left\{ \Phi \left(\text{diagram} \right) + \right. \\ &\quad \left. + R \left[\Phi \left(3 \text{diagram} + 2 \text{diagram} + \text{diagram} \right) \right] \Phi \left(\text{diagram} \right) \right\} \end{aligned}$$

sub-Hopf algebras

- ▶ summing order by order

$$c_k^r = \sum_{|\Gamma|=k, \text{res}(\Gamma)=r} \frac{1}{|\text{Aut}(\Gamma)|} \Gamma \Rightarrow \Delta(c_k^r) = \sum_j \text{Pol}_j(c_m^s) \otimes c_{k-j}^r. \quad (6)$$

sub-Hopf algebras

- ▶ summing order by order

$$c_k^r = \sum_{|\Gamma|=k, \text{res}(\Gamma)=r} \frac{1}{|\text{Aut}(\Gamma)|} \Gamma \Rightarrow \Delta(c_k^r) = \sum_j \text{Pol}_j(c_m^s) \otimes c_{k-j}^r. \quad (6)$$

- ▶ Hochschild closedness

$$X^r = 1 \pm \sum_j c_j^r \alpha^j = 1 \pm \sum_j \alpha^j B_+^{r,j}(X^r Q^j(\alpha)), \quad (7)$$

$$Q^j = \frac{X^v}{\sqrt{\prod_{\text{edges } e \text{ at } v} X^e}}. \text{ Evaluates to invariant charge.}$$

sub-Hopf algebras

- ▶ summing order by order

$$c_k^r = \sum_{|\Gamma|=k, \text{res}(\Gamma)=r} \frac{1}{|Aut(\Gamma)|} \Gamma \Rightarrow \Delta(c_k^r) = \sum_j \text{Pol}_j(c_m^s) \otimes c_{k-j}^r. \quad (6)$$

- ▶ Hochschild closedness

$$X^r = 1 \pm \sum_j c_j^r \alpha^j = 1 \pm \sum_j \alpha^j B_+^{r;j}(X^r Q^j(\alpha)), \quad (7)$$

$$Q^j = \frac{X^v}{\sqrt{\prod_{\text{edges } e \text{ at } v} X^e}}. \text{ Evaluates to invariant charge.}$$

- ▶ $bB_+^{r;j} = 0$.

$$\Delta B_+^{r;j}(X) = B_+^{r;j}(X) \otimes 1 + (id \otimes B_+^{r;j})\Delta(X). \quad (8)$$

Implies locality of counterterms upon application of Feynman rules

$$\Phi B_+^{r;j}(X) = \int d\mu_{r;j} \Phi(X):$$

$$\bar{R}(\Gamma) = m(S_\Phi^R \otimes \Phi P) \Delta B_+^{r;j} = \int d\mu_{r;j} \Phi^R(X). \quad (9)$$

Symmetry

- ▶ Ward and Slavnov–Taylor ids

$$i_k := c_k \bar{\psi} \psi + c_k \bar{\psi} \not{A} \psi \quad (10)$$

span Hopf (co-)ideal I :

$$\Delta(I) \subseteq H \otimes I + I \otimes H. \quad (11)$$

$$\Delta(i_2) = i_2 \otimes 1 + 1 \otimes i_2 + (c_1^{\frac{1}{4}} F^2 + c_1 \bar{\psi} \not{A} \psi + i_1) \otimes i_1 + i_1 \otimes c_1 \bar{\psi} \not{A} \psi.$$

Symmetry

- ▶ Ward and Slavnov–Taylor ids

$$i_k := c_k \bar{\psi} \psi + c_k \bar{\psi} \mathcal{A} \psi \quad (10)$$

span Hopf (co-)ideal I :

$$\Delta(I) \subseteq H \otimes I + I \otimes H. \quad (11)$$

$$\Delta(i_2) = i_2 \otimes 1 + 1 \otimes i_2 + (c_1^{\frac{1}{4}} F^2 + c_1 \bar{\psi} \mathcal{A} \psi + i_1) \otimes i_1 + i_1 \otimes c_1 \bar{\psi} \mathcal{A} \psi.$$

- ▶ Feynman rules vanish on $I \Leftrightarrow$ Feynman rules respect quantized symmetry:

$$\Phi^R : H/I \rightarrow V.$$

Symmetry

- ▶ Ward and Slavnov–Taylor ids

$$i_k := c_k \bar{\psi} \psi + c_k \bar{\psi} \mathcal{A} \psi \quad (10)$$

span Hopf (co-)ideal I :

$$\Delta(I) \subseteq H \otimes I + I \otimes H. \quad (11)$$

$$\Delta(i_2) = i_2 \otimes 1 + 1 \otimes i_2 + (c_1^{\frac{1}{4}} F^2 + c_1 \bar{\psi} \mathcal{A} \psi + i_1) \otimes i_1 + i_1 \otimes c_1 \bar{\psi} \mathcal{A} \psi.$$

- ▶ Feynman rules vanish on $I \Leftrightarrow$ Feynman rules respect quantized symmetry:

$$\Phi^R : H/I \rightarrow V.$$

- ▶ Ideals for Slavnov–Taylor ids generated by equality of renormalized charges, also for the master equation in Batalin–Vilkovisky (see Walter van Suijlekom’s work)

Kinematics and Cohomology

- ▶ Exact co-cycles

$$[B_+^{rj}] = B_+^{rj} + b\phi^{rj} \quad (12)$$

with $\phi^{rj} : H \rightarrow \mathbb{C}$

Kinematics and Cohomology

- ▶ Exact co-cycles

$$[B_+^{rj}] = B_+^{rj} + b\phi^{rj} \quad (12)$$

with $\phi^{rj} : H \rightarrow \mathbb{C}$

- ▶ Variation of momenta

$$G^R(\{g\}, \ln s, \{\Theta\}) = 1 \pm \Phi_{\ln s, \{\Theta\}}^R(X^r(\{g\})) \quad (13)$$

with $X^r = 1 \pm \sum_j g^j B_+^{rj}(X^r Q^j(g))$, $bB_+^{rj} = 0$. Also,

$$G^r = \left[\sum_{j=1}^{\infty} \gamma_j(\{g\}, \{\Theta\}) \ln^j s \right] + \overbrace{G_0^r}^{\text{abelian factor}} \quad (14)$$

Then, for MOM and similar schemes (not MS!):

$$\{\Theta\} \rightarrow \{\Theta'\} \Leftrightarrow B_+^{rj} \rightarrow B_+^{rj} + b\phi^{rj}.$$

$$\Phi_{L_1+L_2, \{\Theta\}}^R = \Phi_{L_1, \{\Theta\}}^R \star \Phi_{L_2, \{\Theta\}}^R.$$

Leading log expansions and the RGE

- ▶ The invariant charge Q^v
For each vertex v , a charge Q^v :

$$Q^v(g) = \frac{X^v(g)}{\prod_e \sqrt{X^e}}, \quad (15)$$

e adjacent to v .

Leading log expansions and the RGE

- ▶ The invariant charge Q^v

For each vertex v , a charge Q^v :

$$Q^v(g) = \frac{X^v(g)}{\prod_e \sqrt{X^e}}, \quad (15)$$

e adjacent to v .

- ▶

$$\left(\partial_L + \beta(g) \partial_g - \sum_{e \text{ adj } r} \gamma_1^e \right) G^r(g, L) = 0 \quad (16)$$

rewrites in terms of the Dykin operator

($\gamma_1^r(g) = S \star Y(X^r(g))$):

$$\gamma_k^r(g) = \frac{1}{k} \left(\gamma_1^r(g) - \sum_{j \in R} s_j \gamma_1^j g \partial_g \right) \gamma_{k-1}^r(g) \quad (17)$$

Ordinary differential equations vs DSE

- ▶ RGE+DSE
the iterated integral structure

$$\Phi^R(B_+^{r;j}(X)) = \int \Phi^R(X) d\mu_{r;j} \quad (18)$$

allows to combine $X^r = 1 \pm \sum_j B_+(X^r Q^j)$ with RGE to

$$\gamma_1^r = P(g) - [\gamma_1^r(g)]^2 + \sum_{j \in R} s_j \gamma_1^j g \partial_g \gamma_1^r(g). \quad (19)$$

Ordinary differential equations vs DSE

- ▶ RGE+DSE
the iterated integral structure

$$\Phi^R(B_+^{r;j}(X)) = \int \Phi^R(X) d\mu_{r;j} \quad (18)$$

allows to combine $X^r = 1 \pm \sum_j B_+(X^r Q^j)$ with RGE to

$$\gamma_1^r = P(g) - [\gamma_1^r(g)]^2 + \sum_{j \in R} s_j \gamma_1^j g \partial_g \gamma_1^r(g). \quad (19)$$

- ▶ massless gauge theories
 $\beta(g) = g\gamma_1(g)/2$ for γ_1 anomalous dim of gauge propagator

$$\gamma_1(g) = \overbrace{P(g)}^{\text{existence assumed}} - \gamma_1(g)(1 - g\partial_g)\gamma_1(g) \quad (20)$$

(Ward Id QED, background field gauge (Abbott) QCD)

Limiting mixed Hodge structures

- ▶ Hopf algebra from flags

$$f := \gamma_1 \subset \gamma_2 \subset \dots \subset \Gamma, \Delta'(\gamma_{i+1}/\gamma_i) = 0 \quad (21)$$

The set of all such flags $F_\Gamma \ni f$ determines Hopf algebra structure, $|F_\Gamma|$ is the length of the flag.

Limiting mixed Hodge structures

- ▶ Hopf algebra from flags

$$f := \gamma_1 \subset \gamma_2 \subset \dots \subset \Gamma, \Delta'(\gamma_{i+1}/\gamma_i) = 0 \quad (21)$$

The set of all such flags $F_\Gamma \ni f$ determines Hopf algebra structure, $|F_\Gamma|$ is the length of the flag.

- ▶ It also determines a column vector $v = v(F_\Gamma)$ and a nilpotent matrix $(N) = (N(|F_\Gamma|))$, $(N)^{k+1} = 0$, $k = \text{corad}(\Gamma)$ such that

$$\lim_{t \rightarrow 0} (e^{-\ln t(N)}) \Phi_R(v(F_\Gamma)) = (c_1^\Gamma(\Theta) \ln s, c_2^\Gamma(\Theta), c_k^\Gamma(\Theta) \ln^k s)^T \quad (22)$$

where k is determined from the co-radical filtration and t is a regulator say for the lower boundary in the parametric representation.

QED

- ▶ sub Hopf algebra for vacuum polarization suffices

QED

- ▶ sub Hopf algebra for vacuum polarization suffices
- ▶ $\gamma_1(x) = P(x) - \gamma_1(x)^2 + \gamma_1(x)x\partial_x\gamma_1(x)$ with $P(x) > 0$

- ▶ sub Hopf algebra for vacuum polarization suffices
- ▶ $\gamma_1(x) = P(x) - \gamma_1(x)^2 + \gamma_1(x)x\partial_x\gamma_1(x)$ with $P(x) > 0$
 $P(x)$ twice

differentiable

$$\gamma_1(x_0) = \gamma_0 > 0$$

different solutions

distinguished by $e^{-\frac{1}{x}}$

behaviour


$$\frac{d\gamma_1}{dx} = \gamma_1 - \gamma_1^2 - P,$$

$$\frac{dx}{dL} = x\gamma_1$$

$$L = \int_{x_0}^{x(L)} \frac{dz}{z\gamma_1(z)}$$

- ▶ **separatrix exists and might have no Landau pole:**

$$D(P) = \int_{x_0}^{\infty} \frac{P(z)dz}{z^3} < \infty, \int_{x_0}^{\infty} \frac{2dz}{z\sqrt{1+4P(z)-1}} < \infty$$



QCD

- ▶ sub Hopf algebra for gluon polarization suffices in background field gauge

QCD

- ▶ sub Hopf algebra for gluon polarization suffices in background field gauge
- ▶ $\gamma_1(g) = P(g) - \gamma_1(g)^2 + \gamma_1(g)g\partial_g\gamma_1(g)$ with $P(g) < 0$

QCD

- ▶ sub Hopf algebra for gluon polarization suffices in background field gauge

- ▶ $\gamma_1(g) = P(g) - \gamma_1(g)^2 + \gamma_1(g)g\partial_g\gamma_1(g)$ with $P(g) < 0$

$P(g)$ twice differentiable
and concave near 0

unique solution which

flows into $(0, 0)$ at large Q^2

$$L = \int_{g_0}^{g(L)} \frac{dz}{z\gamma_1(z)} \rightarrow$$

$$L_\Lambda = - \int_{g(L_\Lambda)}^{\infty} \frac{dz}{z\gamma_1(z)},$$

$$L_\Lambda = \ln Q^2 / \Lambda_{QCD}$$

$$f_{disp}(Q^2) = \int_0^\infty \frac{\Im(f(\sigma))d\sigma}{\sigma + Q^2 - i\eta}$$

and ODE

- ▶ separatrix exists and gives asymptotic free solution, with finite mass gap for inverse propagator iff $\gamma_1(x) < -1$ for some $x > 0$.

$|D(P)| < \infty \rightarrow \gamma_1(x) \sim sx, x \rightarrow \infty$. That allows for dispersive methods as introduced by Shirkov et.al. in field theory.

